

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

Authors: Dena A. Hashim, Alaa Sh. Abdulbari, Noor M. Ali, Alaa Hussein J. Al-Qaisi, Ahmed Rahmah Abu-Raghif, Hayder Ridha-Salman

Article type: Original Article

Received: 28 February 2025

Accepted: 5 June 2025

Published online: 15 August 2025

eISSN: 2544-1361

Eur J Clin Exp Med

doi: 10.15584/ejcem.2025.4.3

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our authors we are providing this early version of the manuscript. The manuscript will undergo copyediting and typesetting. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Exploring the role of B complex vitamins in reproductive health – valuable insights and unresolved issues regarding premature ovarian failure

Dena A. Hashim ¹, Alaa Sh. Abdulbari ², Noor M. Ali ¹, Alaa Hussein J. Al-Qaisi ¹, Ahmed Rahmah Abu-Raghif ³, Hayder Ridha-Salman ⁴

¹ Department of Chemistry, College of Sciences, Al-Nahrain University, AL-Jadriya, Baghdad, Iraq

² Department of Applied Chemistry, College of Applied Sciences, University of Technology-Iraq, Baghdad, Iraq

³ Department of Pharmacology, College of Medicine, Al-Nahrain University, Baghdad, Iraq

⁴ Al-Mustaqbal University, College of Pharmacy, Hillah, Babylon, Iraq

Corresponding author: Hayder Ridha-Salman, e-mail: hayder80.ridha@gmail.com

ORCID

DAH: <https://orcid.org/0000-0002-3505-2711>

ASA: <https://orcid.org/0000-0002-0394-861X>

NMA: <https://orcid.org/0000-0001-7503-6049>

AHJA-Q: <https://orcid.org/0000-0002-6576-0145>

ARA-R: <https://orcid.org/0000-0003-4514-3892>

HR-S: <https://orcid.org/0000-0003-1594-4883>

ABSTRACT

Introduction and aim. Premature ovarian failure (POF) is a condition distinguished by a decline in ovarian follicles in women under 40 leading to a disruption in the normal menstrual cycle and fertility. Vitamin B₆ (pyridoxine) and vitamin B₁₂ (cyanocobalamin) play an important role in ovarian health. The aim was to investigate the effects of vitamins B₆ and B₁₂, for predicting POF via the association of these vitamins with other biochemical parameters.

Material and methods. A total of 80 participants with age ≤ 40 years, 40 POF women, and 40 fertile control women were recruited in this study from November 2023 to March 2024. Serum levels of prolactin (PRO), thyroid hormones, estradiol (E-2), progesterone, follicle stimulating hormone (FSH), luteinizing hormone (LH), 17-alpha hydroxylase, and inhibin alpha (INH- α) were measured using the ELISA technique while vitamins B₆ and B₁₂ were measured by HPLC.

Results. Serum levels of reproductive hormones (LH, E2, progesterone, and testosterone), vitamin B₆, B-

$_{12}$, T_3 , T_4 , 17-alpha hydroxylase, and INH- α were markedly reduced in POF patients, while levels of FSH, TSH, and PRO were significantly elevated compared to healthy controls.

Conclusion. This study highlights the critical role of vitamins B- $_6$ and B- $_{12}$ in the pathophysiology of POF, suggesting their potential as biomarkers for diagnosis and prognosis.

Keywords. 17-alpha-hydroxylase, inhibin-alpha, premature ovarian failure, vitamin B- $_{12}$, vitamin B- $_6$

Introduction

Premature ovarian failure (POF) is a condition that affects women's reproductive health, with its occurrence becoming more prevalent in recent years.¹ POF, often referred to as premature menopause, occurs when a woman experiences cessation of ovarian activity and amenorrhea before turning 40, along with the serum FSH level that rise to the menopausal range. This condition can result in infertility and can cause both physical and emotional challenges.²⁻⁴ Women with POF face a higher risk of psychological disorders, cardiovascular problems, osteoporosis, autoimmune diseases, cognitive impairment, and infection of the urinary and reproductive systems compared to those without the condition.⁵⁻⁷ Ovarian follicles are composed of three types of cells: granulosa cells, oocytes, and theca cell. Granulosa cells and oocytes contain follicle-stimulating hormone (FSH) and luteinizing hormone (LH) receptors, which play a critical role in follicular growth and development.⁸⁻¹⁰ Folliculogenesis is a structured process involving the maturation of follicles in preparation for ovulation.¹¹⁻¹³ However, this process is disrupted in cases of POF.^{1,14,15} Biochemically, POF is defined by reduced levels of gonadal hormones, such as estrogens and inhibins, accompanied by elevated gonadotropins (LH and FSH), a state referred to as hypergonadotropic amenorrhea. This increase in FSH is typically more pronounced than that of LH, with FSH levels exceeding 30 IU/L serving as a diagnostic indicator of ovarian failure.¹⁶⁻¹⁹

Genetic, immunological, metabolic, viral, and medicinal factors all contribute to the etiology of POF.^{2,20-23} B complex, as water-soluble vitamin, is a critical compound for overall cellular metabolism.²⁴ Pyridoxine (B- $_6$) plays an essential role in the regulation of thyroid and sex hormones, the promotion of ovarian maturation, and the production of vital female hormones such as estrogen and progesterone. It is believed to regulate the activity of various steroid hormone receptors at the transcriptional level.²⁵ Cobalamin (B- $_{12}$) is a corrin-based compound containing a corrin ring and plays a critical role in human somatic cell metabolism.²⁶ B- $_{12}$ is essential for nervous and reproductive systems, supporting normal RBC formation, growth, synthesis of DNA, as well as amino acid, fatty acid, and cellular energy metabolism.²⁷⁻²⁹ Once inside the cell, B- $_{12}$ is converted into two active forms: adenosylcobalamin and methylcobalamin. Methylcobalamin functions as a coenzyme for methionine synthase, which plays vital role for metabolism of methionine-homocysteine, commonly referred to as the cycle of methionine. B- $_{12}$, a key player in one carbon metabolism, is a complex metabolic factor essential for human reproductive health.³⁰ Vitamins, particularly B- $_{12}$, are crucial for reproductive health, promoting organ function and mitigating oxidative

stress, pro-inflammatory cytokines, and circulating homocysteine levels.³¹⁻³³ Homocysteine acts as a metabolic linker that hinders the synthesis of intracellular antioxidants. It inhibits key antioxidant enzymes, such as superoxide dismutase and glutathione peroxidase.³⁴ A subclinical deficiency of B-12 impairs the conversion of excess homocysteine to methionine, leading to an increase in homocysteine levels. This, in turn, promotes the production of H₂O₂, which contributes to the accumulation of reactive oxygen species (ROS) accumulation.³⁵⁻³⁷ Oxidative stress, caused by an imbalance between the generation of free radicals and antioxidant levels, can cause detrimental transformation of mitochondrial DNA and protein damage, inevitably provoking cell death.³⁸⁻⁴¹ Methionine serves as a key precursor for S-adenosyl-methionine, an avital compound required for the methylation of phospholipid, neurotransmitters, amines, DNA, RNA and myelin basic protein.^{42,43} A reduction in S-adenosyl-methionine levels can disrupt DNA methylation, potentially altering fetal metabolic programming and increasing susceptibility to non-communicable diseases later in life.⁴⁴ Consequently, B-12 deficiency may lead to homocysteine buildup and reduced circulation methionine, impacting reproduction, as well as protein and DNA methylation.^{45,46} The disruptions in redox balance, DNA methylation, and mitochondrial function are particularly detrimental to ovarian tissue, which is highly sensitive to oxidative stress. Granulosa cells, essential for follicle maturation, are especially vulnerable. Experimental studies have shown that elevated homocysteine levels promote apoptosis in granulosa cells, contributing to follicular atresia. Kaplan and Türk showed that B-12 administration reduced histological markers of tissue damage in rat ovaries by reducing oxidative stress.³⁴ Additionally, Deniz et al. revealed that treatment with B-12 was effective in decreasing inflammation and oxidative stress caused by ovarian ischemia-reperfusion by downregulating the TLR-4/NF-κB signaling pathway, which is associated with inflammation.⁴⁷

The enzyme 17-alpha hydroxylase is expressed in the adrenal glands and gonads, serving as a key branching point in the synthesis of cortisol and sex hormones. This enzyme facilitates two distinct steps in steroidogenesis. Individuals with deficiency in this enzyme commonly present with hypertension, hypokalemia, amenorrhea and absent of secondary sexual characteristics.^{48,49} Inhibin-alpha (INH-α) Plays a role in regulating follicle development by responding to increased levels of FSH in gonads, which in turn triggers negative feedback to the anterior pituitary, reducing circulating FSH levels.⁵⁰⁻⁵² The interaction between vitamins of the B complex, particularly B-6 and Cobalamin (B-12), and the risk of POF remain intricate and not yet fully understood.

Aim

This study aims to investigate the correlation between vitamin B-6 and B-12 levels and POF. More research is required to elucidate their roles and potential therapeutic applications.

Material and methods

Subjects

In this study, 80 participants aged 40 years or under were split into two groups: group one consisted of 40 women with POF, whereas group two included forty healthy fertile women as a control group. From November 2023 to March 2024, POF patients were collected from the Higher Institute of Infertility Diagnosis and Assisted Reproductive Technologies located in Baghdad, Iraq. Women with an unclear etiology of POF, autoimmune diseases, and pelvic surgery were excluded. The research was approved by the University of Technology-Iraq Bioethics Committee on Scientific Research, approval no. BCSR11, on March 24, 2024.

Methods

Blood samples were collected from two groups: the first group (healthy control) on the second day of their menstrual cycle and the second group (POF group) under conditions appropriate to their health status. For each participant, 5 mL of peripheral blood was drawn and centrifuged for 5 minutes at room temperature.

ELISA methods technique

The serum obtained was analyzed for hormone levels including FSH, LH, thyroid hormones, prolactin (PRO), progesterone, estradiol (E-2) and testosterone, 17-alpha hydroxylase and INH- α using the ELISA technique (CUSABIO BIOTECH COM), E-2, INH- α , and progesterone measured used competitive inhibition ELAZA kits, the microtiter plate included in the kits is pre-coated with goat anti-rabbit antibody, standards and samples were placed in designated wells along with specific antibody for these hormones and horseradish peroxidase (HRP) conjugated antigen.⁵³⁻⁵⁵ A competitive reaction occurs between HRP labeled to target hormones and the unlabeled form of hormone for antibody binding. A substrate solution is then introduced, leading to color development inversely proportional to hormone concentration in the sample.⁵⁶⁻⁵⁹ The reaction halted and the color intensity is measured, while the other tests was measured by used quantitative sandwich ELAZA kits, the microplate included in the kits is pre-coated with antibody specific for target hormone, Standards and samples were added to wells along horseradish peroxidase (HRP)-conjugated antibody targeting hormone. After washing to remove unbound reagents, a substrate solution is introduced, resulting in color development inversely proportional to the hormone concentration bound in the initial step.⁶⁰⁻⁶² The reaction is stopped and the color intensity is measured.

HPLC technique

Vitamin B-6 and B-12 concentrations were measured using the HPLC technique, the HPLC Autosampler SyKNM from Germany S5200 pump model S1122 Column oven S4011, and a UV detector S3240 set at wavelength 230 nm with a C-18 column., using acetonitrile (A) and NaH₂PO₄ (B), with a gradient of 100%

(B) at 3 minutes, 80% (B) at 6 minutes, and 60% (B) at 15 minutes. The flow rate remained constant at 1.0 mL/min.^{40,63,64} Standard vitamins B-6, B-12 were sourced from Sigma. Creating a standard solution began with dissolving 10 mg of each of vitamins B-6 and B-12 in methanol to form a 100 parts per million (ppm) stock solution, diluted to 100 mL, then the stock solution was further diluted to prepare other standard solutions by subsequent dilution of the stock solutions.⁶⁵⁻⁶⁷

Post hoc power analysis

The results of the post hoc power analysis indicate that the study had a very high ability to detect differences in testosterone levels between the control and POF groups. The mean testosterone level in the control group was 0.47 with a standard deviation of 0.22, while the mean in the POF group was 0.09 with a standard deviation of 0.02. The effect size, calculated using Cohen's d, showed a very large difference between the two groups, reaching 2.55, which reflects a significant variation in testosterone levels. The sample size consisted of 40 individuals in the control group and 40 in the POF group, further strengthening the reliability of the study. The post hoc power reached 1.0 (100%), which means that the study was fully capable of detecting any real effect if it existed, which is a strong indicator of the statistical reliability of the results.




Fig. 1. Post hoc power analysis curve: effect of sample size on statistical power

The post hoc power analysis Figure 1 shows that the study has a very high power to detect differences between the two groups, with the statistical power remaining close to 1.0 (100%) even with small sample sizes. The horizontal axis represents the sample size per group, while the vertical axis reflects the level of statistical power. The orange line indicates how the power remains consistently high across different sample sizes. On the contrary, the red dashed line represents the minimum acceptable power level of 0.8, commonly used as a benchmark for detecting real effects in scientific studies. This result means that the survey can detect any actual differences between the groups, due to the enormous effect size (Cohen's $d = 2.55$), making the likelihood of a Type II error (failing to detect an actual effect) almost nonexistent.

Statistical analysis

All statistical analyses in this study were conducted using statistical Package for Social Science (SPSS) version 29.0 windows (SPSS, Inc., Chicago, IL USA). Descriptive statistics were expressed as mean $\pm$ standard deviation (SD). Differences between mean values were evaluated using Student's t test, with significance threshold set at $p < 0.05$, while $p > 0.05$ was considered nonsignificant.

Results

Table 1 presents the biochemical profiles of the studied groups, highlighting highly significant differences ($p < 0.001$) between the POF group and the control. Serum levels of reproductive hormones LH, E2, progesterone, and testosterone were markedly reduced ($p < 0.001$) in POF patients, while levels of FSH and PRO were significantly elevated compared to healthy controls. Furthermore, thyroid hormones (T3 and T4) showed a significant decrease ($p < 0.001$) in the POF group, accompanied by an increase in TSH levels compared to the control group. Table 2 and Figure 1 highlight significant differences in levels of B₁₂, B₆, and INH- α between the two studied groups: POF and healthy controls. These levels were markedly lower in POF patients compared to the control group.



Table 1. The mean ($\pm$ SD) of age and hormones for the control and POF groups

Characteristic	Mean $\pm$ SD		95% CI	p
	Control group (n=40)	POF group (n=40)		
Age (year)	32.07 $\pm$ 5.370	32.92 $\pm$ 4.26	-3.00–1.30	0.43
FSH (mIU/mL)	3.40 $\pm$ 1.10	38.64 $\pm$ 11.59	-38.95– 31.51	<0.001
LH (mIU/mL)	3.67 $\pm$ 1.64	0.83 $\pm$ 0.40	2.30–3.38	<0.001
E-2 (pg/mL)	160.31 $\pm$ 38.76	10.93 $\pm$ 2.31	136.9–	<0.001

			161.76	
Progesterone (ng/mL)	2.27 ±2.03	0.06±0.062	1.56–2.86	<0.001
PRO (ng/mL)	14.62±4.09	38.25±7.25	-26.26– 20.99	<0.001
Testosterone (ng/mL)	0.47±0.22	0.09 ±0.022	0.31–0.45	<0.001
TSH (mIU/L)	2.75±1.11	8.18±1.87	0.34–6.1	<0.001
T ₃ (nmol/L)	1.46±0.36	0.68±0.09	0.05–0.66	<0.001
T ₄ (nmol/L)	122.99±23.82	40.49±15.62	4.50–73.50	<0.001

Table 2. The mean (±SD) of vitamin B-6, vitamin B-12, 17-alpha hydroxylase, and inhibin-α for the control and POF groups

Characteristic	(Mean±SD)		95 % CI	p
	Control group (n=40)	POF group (n=40)		
Vitamin B-6 (ng/mL)	40.68±5.75	15.17±3.81	23.32–27.68	<0.001
Vitamin B-12 (ng/mL)	30.68±5.75	5.45±3.42	23.11–27.37	<0.001
17-alpha hydroxylase (ng/dL)	50.68±47.17	0.032±0.018	35.56–65.73	<0.001
INH-α (ng/mL)	29.12±9.11	3.26±1.28	22.91–28.80	<0.001






Fig. 2. Mean distribution of biochemical parameters in POF patients group and control group

Discussion

The eight water-soluble vitamins that make up the B-vitamin complex cooperate as coenzymes in an array of anabolic and catabolic enzyme activities to play vital roles in cellular activity.^{24,32} Vitamin B₆ is necessary for the synthesis and metabolism of various hormones, including estrogen and progesterone, which is crucial for menstrual regularity, fertility, and overall reproductive health.⁶⁸ Several studies evaluated B-vitamin circulating levels in the context of neuroimmune diseases.⁶⁹⁻⁷¹ The present study revealed the evaluation of the effect of B₆ and B₁₂ vitamins in POF patients and suggests that they are related to POF as well as hormonal dysregulation that occurs with ovarian disorders.

The current study demonstrated that B₆ and B₁₂ vitamins were significantly lower in POF patients compared to healthy control. These results align with studies conducted by Brilliant et al.⁷² and Kamrul-Hasan et al.⁷³, which also reported deficiencies of these vitamins in female reproductive disease, highlighting their potential role in ovarian health.

We did not account for dietary intake, supplementation, or lifestyle factors that could influence these levels. However, it is important to note that none of the participants in our study took vitamin supplements minimizing potential confounding effects for supplementation.

Antioxidant micronutrients, such as vitamin and minerals, are vital for female reproductive health as they help manage oxidative stress and facilitate essential cellular function.⁷⁴⁻⁷⁶ Recent studies indicate that B vitamins could influence ovarian health and function. Specifically, B₁₂ can help mitigate ovarian ischemia-reperfusion injury by reducing inflammation and oxidative stress.⁴⁷ Similarly, B₆, which is critical for reproduction, affects sulfur and seleno amino acid metabolism and may influence ovarian health and

embryo development under oxidative stress.⁷⁷ Although not explicitly related to POF, vitamin B-6 has been shown to lower serum PRO levels in hyperprolactinemic patients, a condition relevant to reproductive health.⁷⁸ These findings hint at a relationship between vitamins and ovarian function, but more research is required to confirm these links and explore their therapeutic potential in POF.

17-alpha hydroxylase enzyme was significantly lower in POF group compared to the control group which agrees with certain theory reported that a lower level of 17-alpha hydroxylase enzyme alters ovarian functions and the production of sex hormones, as this enzyme is responsible for converting progesterone into cortisol.⁷⁹ Women with 17-alpha hydroxylase deficiency often experience infertility and related complications due to disruptions in the steroidogenic pathway. These abnormalities lead to impaired levels of androstenediones, 17-hydroxyprogesterone, and E-2.⁸⁰ Weak estrogen feedback to the pituitary may result in mildly elevated FSH and LH levels, increasing the risk of ovarian cysts, particularly multiple luteinizing ovarian cysts. Furthermore, increasing PRO levels may be due to increased GnRH activity. Another common reproductive endocrine manifestation in women with 17-alpha hydroxylase deficiency is deduced testosterone levels. However, in females, low testosterone has minimal impact on reproductive development and fertility. However, this characteristic can serve as a clinical indicator, emphasizing the need to consider 17-alpha hydroxylase deficiency in cases where pubic hair is absent or absent during gynecological examination. In general, partial 17-alpha hydroxylase deficiency presents with endocrine abnormalities that correspond to the degree of enzymatic dysfunction.⁸¹

The current study demonstrated a significant decrease in the INH- α ($p < 0.001$) in POF group compared to the control group. Ovarian follicle granulosa cells are the main producers of INH- α . Blocks the ability of the pituitary gland to secrete FSH, which is its primary role. INH- α is produced by mature and developing follicles during a normal menstrual cycle. This hormone works with the pituitary gland to decrease FSH levels.⁸² In accordance with this theory, the reduced level of INH- α level in patients with POF increases the release of FSH by eliminating the inhibitory feedback of the pituitary gland. Several studies have reported a lower serum inhibin in individuals with POF.^{83,84} This deficiency is believed to contribute to disease development, as it leads to increased activation activity. In turn, activation stimulates FSH secretion, accelerating folliculogenesis and speeding up ovarian follicle depletion, which ultimately results in early menopause.⁸⁵ Genetic variation in genes responsible for inhibin production has been associated with its reduced serum levels, which may amplify activation effects due to diminished inhibin secretion.⁸⁵ Previous study identified INH- α (rs12720062 G / A) and showed genotype GG and allele G as potential risk factors for POF, while genotype GA and allele A appear productive. However, genotype AA does not seem to have a significant influence on disease susceptibility.⁸⁶ The reduced levels of 17-alpha hydroxylase and INH- α observed in POF patients' highlights their critical roles in maintaining ovarian function. While direct evidence linking vitamin B-6 and B-12 to the synthesis of these proteins is limited, their fundamental roles in enzymatic reactions and gene expression suggest that deficiency could adversely affect ovarian health.

More research is necessary to elucidate these relationships and determine whether supplementation could serve as a therapeutic strategy for POF. Given the observed association between vitamin B₋₁₂ and B₋₆ levels and POF, it is essential to explore whether correcting such deficiencies by targeted supplementation might influence disease progression, severity of symptoms, or fertility outcomes of affected women. Randomized controlled trials are recommended to evaluate this hypothesis.

Reduced PRO, thyroid hormones (T₃, T₄), LH, E2, and progesterone in patients with POF due to ovarian dysfunction and altered hormonal functions can all contribute to the development of POF or exacerbate its symptoms. Support this theory; a study reported that infertility in women is indicated by the significant positive correlation TSH has with PRO, FSH, and LH. When diagnosing infertility, these hormonal assessments enable a standard etiological approach.^{87,88}

Our participants were recruited from Baghdad, which may limit the generalizability of the findings. Future studies should consider the possible impact of ethnic and genetic diversity on vitamin metabolism and ovarian health.

Study limitations and recommendations

This study provides a cross-sectional initial snapshot of biochemical parameters in POF patients and does not establish causality. A key limitation of this study is the relatively small sample size (80 participants), which may affect the statistical power and generalizability of the results. Although our findings provide valuable information on the possible association between B₋₆ and B₋₁₂ and POF, larger-scale longitudinal studies involving more diverse populations to validate these observations. Further research should incorporate dietary assessment, supplement intake, physical activity, and genetic analysis to provide a more comprehensive understanding of the factors that influence vitamin levels in POF. These unmeasured variables could introduce bias and limit the generalizability of the findings. Another major drawback is the lack of population diversity; more research involving participants from multiple centers is necessary to improve external validity in examining possible environment-gene interactions that can affect POF risk. However, the existing literature suggests that adequate levels of vitamin B are necessary for women's health, especially ovarian function, but also that further investigated studies are required to determine their therapeutic role in POF management.

Conclusion

The study findings revealed a significant deficiency in vitamin B₋₆, B₋₁₂, INH- α , and 17- α hydroxylase levels in patients with POF. These results suggest that POF is associated with disturbances in vitamin B₋₆ and B₋₁₂, as well as dysregulation of INH- α and 17- α hydroxylase production. These biochemical abnormalities may contribute to the pathogenesis and clinical manifestations of POF. More research is needed to elucidate the underlying mechanisms and explore potential therapeutic interventions that target

these deficiencies, aiming to improve the diagnosis and management of POF patients.

Declarations

Funding

The authors received no financial support from economic organizations.

Author contributions

Conceptualization, D.A.H. and A.S.A.; Methodology, N.M.A., A.H.J.A. and A.R.A.; Software, H.R.-S.; Validation, H.R.-S., A.R.A. and A.H.J.A.; Formal Analysis, D.A.H.; Investigation, N.M.A. and A.H.J.A.; Resources, D.A.H. and A.S.A.; Data Curation, A.H.J.A.; Writing – Original Draft Preparation, A.S.A.; Writing – Review & Editing, A.R.A. and H.R.-S.; Visualization, N.M.A.; Supervision, A.R.A. and H.R.-S.; Project Administration, N.M.A.; Funding Acquisition, D.A.H. and A.S.A.

Conflicts of interest

The authors declare that they have no conflict of interest.

Data availability

The data supporting the findings of this study are available from the corresponding author upon reasonable request.

Ethics approval

The study protocol was approved by the University of Technology-Iraq Bioethics Committee (approval no.: BCSR11) on 24 March 2024. All procedures were in accordance with internationally recognized, ethical standards, particularly those in the Declaration of Helsinki.

References

1. Shareghi-Oskoue O, Aghebati-Maleki L, Yousefi M. Transplantation of human umbilical cord mesenchymal stem cells to treat premature ovarian failure. *Stem Cell Res Ther.* 2021;12(1):454. doi: 10.1186/s13287-021-02529-w
2. Bai X, Wang S. Signaling pathway intervention in premature ovarian failure. *Front Med (Lausanne).* 2022;9:999440. doi: 10.3389/fmed.2022.999440
3. Al-Yasiry RZ, Jawad MA, Abbood MS. Association of Methylenetetrahydrofolate Reductase Gene Polymorphism C677T with Oocyte Number and Embryo Quality in Iraqi Infertile Women Undergoing Intracytoplasmic Sperm Injection. Article. *Al-Rafidain Journal of Medical Sciences.* 2024;6(1):39-45. doi: 10.54133/ajms.v6i1.421

4. Kadim Al-Tuma AM, Hussain SS, Hassan MF. Treatment with Metformin Can Improve Insulin Resistance and Lipid Profile in Non-Diabetic Iraqi Women with Polycystic Ovary Syndrome. *Medical Journal of Babylon*. 2024;21(2):S185-S188. doi: 10.4103/MJBI.MJBI_913_923
5. Conte B, Del Mastro L. Gonadotropin-releasing hormone analogues for the prevention of chemotherapy-induced premature ovarian failure in breast cancer patients. *Minerva Ginecol*. 2017;69(4):350-356.
6. Hussein Mohsin HA, Jwad MA, Mossa HA. Is There a Correlation between Serum SFRP5 and Wnt5a Proteins and Insulin Resistance in Iraqi Infertile Females Undergoing ICSI? *Al-Anbar Med J*. 2023;19(2):155-161.
7. Alhussien ZA, Mossa HAL, Abood MS. The Effect of L-carnitine on Apoptotic Markers (Annexin V and Clusterin) in Polycystic Ovarian Syndrome Women undergoing ICSI. Article. *Int. J Drug Deliv Technol*. 2022;12(4):1682-1686. doi: 10.25258/ijddt.12.4.33
8. Sheikhsari G, Aghebati-Maleki L, Nouri M, Jadidi-Niaragh F, Yousefi M. Current approaches for the treatment of premature ovarian failure with stem cell therapy. *Biomed Pharmacother*. 2018;102:254-262.
9. Dawood SA, Jwad MA, Mossa HAL. The impact of serum and follicular fluid irisin on oocyte and embryonic characteristics in infertile women undergoing ICSI according to BMI. *Al-Rafidain Journal of Medical Sciences*. 2023;5:211-217.
10. Ali ZH, Mossa HAL, Abood MS. Evaluation of Endometrial Receptivity and Pregnancy Outcome in view of Assessment of Serum and Follicular Fluid Level of CD44 Following Administration of Hyaluronic Acid to Infertile Women. Article. *Pharmacologyonline*. 2021;3:536-544.
11. Abunaila RSH, Al-Anbari LA, Abbood MS. The effects of adding gonadotropin-releasing hormone antagonist on cycle characteristics and pregnancy rate in stimulated intrauterine insemination (IUI) cycle. *IJRPS*. 2020;11(3):3053-3060. doi: 10.26452/ijrps.v11i3.2407
12. Herez SH, Jawad MA, Abbood MS. Effect of Bisphenol A Level in Follicular Fluid on ICSI Outcome. *J Pharm Negat Results*. 2022;13(4):370-376.
13. Mohsin HAH, Reshan RG, Jawad MA. Comparison between Effects of Ketamine and Remifentanyl Used During Oocyte Retrieval on ICSI Outcome. *Syst Rev Pharm*. 2020;11(4):621-628.
14. Shihab EM, Ahmed HN, Ahmed IT, Abood MN. The story of oral contraceptive users and bacterial vaginosis.... Is it real?. *Int J Pharm Res*. 2020;12:1809-1818. doi: 10.31838/ijpr/2020.SP1.277
15. Sabeeh Hussain S, Majeed Rustum H. Evaluation of anti-zona pellucida antibodies in serum and follicular fluid for polycystic ovarian women undergoing intracytoplasmic sperm injection programme. Article. *IJRPS*. 2018;9(4):1533-1537. doi: 10.26452/ijrps.v9i4.1716
16. Beck-Peccoz P, Persani L. Premature ovarian failure. *Orphanet J Rare Dis*. 2006;1:1-5.
17. Abed FA, Maroof RE, Al-Nakkash UMA. Comparing the diagnostic accuracy of anti-Müllerian

hormone and follicle stimulating hormone in detecting premature ovarian failure in Iraqi women by ROC analysis. *Rep Biochem Mol Biol*. 2019;8(2):126.

18. Jankowska K. Premature ovarian failure. *Prz Menopauzalny*. 2017;16(2):51-56.

19. Raaf GB, Mohammed AA, Jawad MA. The Comparison of the Effect of Recombinant FSH in Antagonist Protocol on Serum and Follicular Fluid Kisspeptin between PCOS and non-PCOS Infertile Women during ICSI. *IJDDT*. 2022;12(1):33-38.

20. Jawad HH, Al-Anbari L, Abbood M. Assessment of the relationship of basal serum Anti-Mullerian hormone levels and maternal age with pregnancy out come in patients undergoing ICSI. *Ann Trop Med Public Health*. 2020;23:11-20.

21. Al-Anbari LA, Jwad MA, Kadhim AAM, Abbood MS, Ridha-Salman H, Al-Athari AJH. Oral intake of *Salvia officinalis* improves intrauterine insemination outcomes in women with polycystic ovary syndrome. *Ginekologia i Poloznictwo*. 2025;20:1-5.

22. Al-yasiry RZ, Jwad MA, Hasan MF, Alsayigh HA. How obesity affects female fertility. *Medical Journal of Babylon*. 2022;19(2):111-114.

23. Abdullah MA, Abdulhamed WA, Abood MS. Effects of Vaginal Sildenafil Citrate on Ovarian Blood Flow and Endometrial Thickness in Infertile Women Undergoing Intra-uterine Insemination. Article. *Int. J Drug Deliv Technol*. 2021;11(4):1450-1453. doi: 10.25258/ijddt.11.4.56

24. Mary Hanna EJ, Van Nguyen, Jeremy Clay. B Vitamins: Functions and Uses in Medicine. *Perm J*. 2022;26(2):89-97. doi: 10.7812/TPP/21.204

25. Hsu JM. Interrelations between vitamin B6 and hormones. *Vitamins & Hormones*. 1964;21:113-134.

26. Obeid R, Heil SG, Verhoeven MMA, van den Heuvel E, de Groot L, Eussen S. Vitamin B12 Intake From Animal Foods, Biomarkers, and Health Aspects. *Front Nutr*. 2019;6:93. doi: 10.3389/fnut.2019.00093

27. Ge Y, Zadeh M, Mohamadzadeh M. Vitamin B12 Regulates the Transcriptional, Metabolic, and Epigenetic Programing in Human Ileal Epithelial Cells. *Nutrients*. 2022;14(14). doi: 10.3390/nu14142825

28. Fakhri SA, Salman HR, Shuubber ZI. Effect of maternal B12 on the child weight. *Bulletin of National Institute of Health Sciences*. 140(3):2405-2411.

29. Khan MG, Hussain SHA, Alkhayl FFA, Ahsan M, Ridha-Salman H. Cannabinoids in neuropathic pain treatment: pharmacological insights and clinical outcomes from recent trials. *Naunyn Schmiedebergs Arch Pharmacol*. 2025. doi: 10.1007/s00210-025-04134-7

30. Wang X, Fu K, Zhang Y, Jiang S, Zhang C. Does Vitamin B12 Influence Clinical Pregnancy Outcome in the follicular-phase GnRH agonist protocol? 2023. doi: 10.21203/rs.3.rs-2904686/v1

31. Chan W, Almasieh M, Catrinescu MM, Levin LA. Cobalamin-Associated Superoxide Scavenging in Neuronal Cells Is a Potential Mechanism for Vitamin B(12)-Deprivation Optic Neuropathy. *Am J Pathol*.

2018;188(1):160-172. doi: 10.1016/j.ajpath.2017.08.032

32. Shakoor H, Feehan J, Mikkelsen K, et al. Be well: A potential role for vitamin B in COVID-19. *Maturitas*. 2021;144:108-111. doi: 10.1016/j.maturitas.2020.08.007

33. Abu-Raghif AR, Alkazzaz AM, Fadheel QJ. A comparative study of the effect of thyme and Calcium with Vitamin D3 in treatment of postmenopausal women with osteoporosis. *Res J Pharm Biol Chem Sci*. 2016;9(5):260-268.

34. Kaplan S, Türk A. Effects of vitamin B12 on rat ovary with ischemia-reperfusion injury. *Biotech Histochem*. 2022;97(4):284-289. doi: 10.1080/10520295.2021.1961863

35. Silverstein WK, Cheung MC, Lin Y. Vitamin B(12) deficiency. *CMAJ*. 2022;194(24):E843. doi: 10.1503/cmaj.220306

36. Rzepka Z, Respondek M, Rok J, et al. Vitamin B(12) Deficiency Induces Imbalance in Melanocytes Homeostasis-A Cellular Basis of Hypocobalaminemia Pigmentary Manifestations. *Int J Mol Sci*. 2018;19(9):2845. doi:10.3390/ijms19092845

37. Sharquie KE, Turki KM, Abu-Raghif AR, Al-Geboori HG. Oxidative stress in patients with premature hair grayness. Article. *Saudi M J*. 2005;26(8):1310-1311.

38. Khaleel BJ, Ridha-Salman H, Kadhim HM, Hassan OM, Kubba A, Sahib HB. Anti-angiogenic and anti-oxidant effects of 2-NTI indole derivative vs. suramin in ex vivo, in vivo, and in vitro studies. *Cytotechnology*. 2025;77(1):38. doi: 10.1007/s10616-024-00701-7

39. Khaleel BJ, Ridha-Salman H, Kadhim HM, Hassan OM, Kubba A, Sahib HB. Inhibitory effects of carbonylhydrazide indole derivative on micro-blood vessel growth using ex vivo, in vivo, and in vitro assays. *In Vitro Cell Dev Biol Anim*. 2025. doi: 10.1007/s11626-025-01019-0

40. Yahya YI, Hadi NR, Abu Raghif A, Qassam H, Al Habooby NGS. Role of Iberin as an anti-apoptotic agent on renal ischemia-reperfusion injury in rats. *J Med Life*. 2023;16(6):915-919. doi: 10.25122/jml-2022-0281

41. Aal-Aaboda M, Abu Raghif A, Hadi N. Renoprotective Potential of the Ultra-Pure Lipopolysaccharide from *Rhodobacter Sphaeroides* on Acutely Injured Kidneys in an Animal Model. *Archives of Razi Institute*. 2021;76(6):1755-1764.

42. Froese DS, Fowler B, Baumgartner MR. Vitamin B(12) , folate, and the methionine remethylation cycle-biochemistry, pathways, and regulation. *J Inherit Metab Dis*. Jul 2019;42(4):673-685. doi: 10.1002/jimd.12009

43. Falh A, Hussain, Sraibet A, Muyad. Correlation Study between Level of zinc and lead in Serum and Seminal Plasma in Infertile Patients.. *Ann Trop Med Public Health*. 2020;23(16):231-246. doi: 10.36295/ASRO.2020.231614

44. Randunu RS, Bertolo RF. The Effects of Maternal and Postnatal Dietary Methyl Nutrients on Epigenetic Changes that Lead to Non-Communicable Diseases in Adulthood. *Int J Mol Sci*.

2020;21(9):3290. doi: 10.3390/ijms21093290

45. Akamine K, Mekaru K, Gibo K, et al. Impact of the one-carbon metabolism on oocyte maturation, fertilization, embryo quality, and subsequent pregnancy. *Reprod Med Biol*. Jan 2021;20(1):76-82. doi: 10.1002/rmb2.12354
46. Agbalalah T, Robert FO, Amabebe E. Impact of vitamin B12 on the reproductive health of women with sickle cell disease: a narrative review. *Reprod Fertil*. 2023;4(3):e230015. doi: 10.1530/raf-23-0015
47. Deniz E, Topcu A, Ozturk A, Ozturk SD, Arpa M, Atak M. The effects of vitamin B12 on the TLR-4/NF- κ B signaling pathway in ovarian ischemia-reperfusion injury-related inflammation. *Int Immunopharmacol*. 2022;107:108676. doi: 10.1016/j.intimp.2022.108676
48. Chen H, Chen Y, Mao H, Huang H, Lou X. A novel homozygous CYP17A1 mutation causes partial 17 α -hydroxylase/17, 20-lyase deficiency in 46, XX: a case report and literature review. *Blood Pressure*. 2023;32(1):2195008.
49. Xia Y, Shi P, Xia J, Zhang H, Xu L, Kong X. Novel mutations of the CYP17A1 gene in four Chinese 46, XX cases with partial 17 α -hydroxylase/17, 20-lyase deficiency. *Steroids*. 2021;173:108873.
50. Zhang Y, Chen Z, An C, et al. Effect of active immunization with recombinant-derived goose INH- α , AMH, and PRL fusion protein on broodiness onset and egg production in geese (*Anser cygnoides*). *Poultry Science*. 2021;100(11):101452.
51. Akhtar MF, Wei Q, Zhu H, et al. The role of active immunization against inhibin α -subunit on testicular development, testosterone concentration and relevant genes expressions in testis, hypothalamus and pituitary glands in Yangzhou goose ganders. *Theriogenology*. 2019;128:122-132.
52. Hussain SS, Hassan MF. The effectiveness of using r-HMG+r-FSH vs. R-FSH alone during COS on ICSI outcome. Article. *Ann Trop Med Public Health*. 2019;22(8):S232. doi: 10.36295/ASRO.2019.22089
53. Luty RS, Abbas SF, Haji SA, Ridha-Salman H. Hepatoprotective effect of catechin on rat model of acetaminophen-induced liver injury. *Comp. Clin. Path*. 2025. doi:10.1007/s00580-025-03682-x
54. Attarbashee RK, Hamodat HF, Mammdoh JK, Ridha-Salman H. The Possible effect of Bosentan on the methotrexate-induced salivary gland changes in male rats: histological and Immunohistochemical study. *Toxicology Research*. 2025;14(1):tfaf007. doi: 10.1093/toxres/tfaf007
55. Luty RS, Al-Zubaidy AA, Malik AS, Ridha-Salman H, Abbas AH. Protective effect of orientin on diabetic nephropathy in rat models of high-fat diet and streptozotocin-induced diabetes. *Naunyn Schmiedebergs Arch Pharmacol*. 2025. doi:10.1007/s00210-025-03949-8
56. Raheem AKK, Abu-Raghif AR, Zigam QA. Cilostazol Protects Against Sepsis-Induced Kidney Impairment in a Mice Model. *J Med Chem Scie*. 2023;6(5):1193-1203. doi: 10.26655/jmchemsci.2023.5.25
57. Habbas AH, Abu-Raghif AR, Ridha-Salman H, Hussein MN. Therapeutic effect of bosentan on 2, 4-dinitrochlorobenzene (DNCB)-induced atopic dermatitis mouse model. *Arch Dermatol Res*.

2025;317(1):436. doi: 10.1007/s00403-025-03955-z

58. Abbas AH, Abbood MS, Ridha-Salman H, Fakhri SA, Abbas ZH, Al-Athari AJH. Suppressive effect of topical moxifloxacin on imiquimod-induced model of psoriasis in mice. *Naunyn Schmiedeberg's Arch Pharmacol*. 2025. doi: 10.1007/s00210-025-04317-2
59. Ali BF, Abu-Raghif AR, Ridha-Salman H, Al-Athari AJH. Vildagliptin topical ointment: an effective treatment for imiquimod-induced psoriasis in mice. *J Mol Histol*. 2025;56(3):143. doi: 10.1007/s10735-025-10416-4
60. Kadhim H, Gatea F, Raghif AA, Ali K. Role of Topical Ritodrine Hydrochloride in Experimentally Induced Hypertrophic Scar in Rabbits. *Iraqi J Pharm Sci*. 2022;31(2):260-270. doi: 0.31351/vol31iss2pp260-270
61. Khorsheed SM, Abu-Raghif A, Ridha-Salman H. Alleviative Effects of Combined Topical Melatonin and Rutin on Imiquimod-Induced Psoriasis Mouse Model. 10.3897/pharmacia.71.e128832. *Pharmacia*. 2024;71:1-13. doi: 10.3897/pharmacia.71.e128832
62. Raheem AK, Abu-Raghif AR, Abbas AH, Ridha-Salman H, Oubaid EN. Quercetin mitigates sepsis-induced renal injury via inhibiting inflammatory and oxidative pathways in mice. *J Mol Histol*. 2025;56(3):184. doi: 10.1007/s10735-025-10442-2
63. Sati'Abbas S, Abul-Razak N, Mustafa N, Abd Ali R. Homocysteine, folic acid, vitamin B12 and pyridoxine: effects on vaso-occlusive crisis in sickle cell anemia and sickle-thalassemia. *Iraqi Postgraduate Medical Journal*. 2011;10(4):473-479.
64. Manna MJ, Abu-rahif A, Muhsin HY. The effect of Niclosamide in acetic acid induce colitis: an experimental study. *Prensa méd argent*. 2019;105(5):309-316.
65. Oubaid EN, Abu-Raghif AR, Al-Sudani IM. Phytochemical Screening and Antioxidant Activity of *Uncaria tomentosa* Extract: In Vitro: and: In Vivo: Studies. *M J Babylon*. 2023;20(1):136-142.
66. Hassan Z, Hassan T, Abu-Rghif A. Evaluation the Effectiveness of Phenolic Compound of *Salvia frugida* on Induced Atopic Dermatitis in Experimental Mice. *Iraqi J P Sci*. 2022;31(1):154-166. doi: 10.31351/vol31iss1pp154-166
67. Thammar MR, Sahib HB, Ridha-Salman H. Skin Healing Potential of Bioactive Components From *Lycoperdon lividum* Mushroom Versus β -Sitosterol in Rat Model of Burn Wounds. *Microscopy Research and Technique*. 2025;88:1-53. doi: 10.1002/jemt.24864
68. Zhao Y, Hu S, Zhai W, Wang M, Ran L. [Retracted] Clinical Study of Progesterone Combined with Vitamin B6 in the Treatment of Amenorrhea Endocrine Disorders Caused by Antipsychotics. *Comput Math Methods Med*. 2022;2022(1):2436322.
69. Bucolo C, Maugeri G, Giunta S, D'Agata V, Drago F, Romano GL. Corneal wound healing and nerve regeneration by novel ophthalmic formulations based on cross-linked sodium hyaluronate, taurine, vitamin B6, and vitamin B12. *Front Pharmacol*. 2023;14:1109291.

70. Tothonglor A, Kobutree P, Roumwong A, Jindatip D, Agthong S. Cisplatin-induced alterations in the blood-nerve barrier: effects of combination of vitamin B1, B6 and B12. *Folia Morphologica*. 2023;82(1):53-62.
71. Abdulbari AS, Ali NM, Abu-Raghif AR, Matloob NA, Ridha-Salman H. Impact of azithromycin on specific biochemical markers and sebum composition in acne vulgaris patients. *Arch Dermatol Res*. 2025;317(1):798. doi: 10.1007/s00403-025-04299-4.
72. Brilliant A, Astuti BPK, Joyo EO, Febri RR, Silvana V, Muharam R. Vitamin B3 (niacin), B6, C, and iron intake are associated with the free androgen index, especially in normoandrogenic polycystic ovary syndrome. *J Turk Ger Gynecol Asso*. 2022;23(3):130.
73. Kamrul-Hasan A, Aalpona FTZ. Comparison of Serum Vitamin B12 Levels Among Drug-Naïve and Metformin-Treated Patients With Polycystic Ovary Syndrome. *Cureus*. 2022;14(10):e30447. doi: 10.7759/cureus.30447
74. Mohammed A, Al-Suwaiegh S, Al-Gherair I, et al. The Adjuvant Impacts of Antioxidant Micronutrients on Ovarian Follicle Development, Oocyte Maturation and Embryo Development of Mammalian Species: A Review. *Indian J Anim Res*. 2024;58(10):1651-1657.
75. Oudah NN, Abdulbari AS, Hameed N. Serum Electrolytes Levels in Third Trimester Pregnant Women: A Case-Control Study in Diyala Province in Iraq. *Egyptian J Chem*. 2022;65(8):31-34.
76. Shihab EM, Kadhim HM. The Impact of Carvedilol on Organ Index, Inflammatory Mediators, Oxidative Stress Parameters and Skin Markers in D-Galactose-Induced Aging Mice. *Int J Drug Del Tech*. 2023;13(3):1017-1023. doi: 10.25258/ijddt.13.3.41
77. Dalto DB, Matte J-J. Nutrigenomic aspects of dietary pyridoxine (vitamin B6) and selenium interaction and their implications in reproduction. *Molecular Nutrition*. 2020:131-151.
78. Witwit SJ. The role of vitamin B6 in reducing serum prolactin in comparison to cabergoline. *IJDFR*. 2019;10(1):108-113.
79. Şimşek Y, Polat S, Riepe FG, et al. 17-Alpha Hydroxylase Deficiency: A Rare Case of Primary Amenorrhea and Hypertension. *Turkish Journal of Endocrinology & Metabolism*. 2014;18(4):137-139.
80. He Z, Ma Y, Li L, et al. Osteopontin and Integrin $\alpha\beta3$ Expression during the Implantation Window in IVF Patients with Elevated Serum Progesterone and Oestradiol Level. *Geburtshilfe Frauenheilkd*. 2016;76(6):709-717. doi: 10.1055/s-0041-111697
81. Du X, Jia Q, Wu S, Wang B, Guan Y. Successful live birth in women with partial 17 α -hydroxylase deficiency: report of two cases. *Reproductive BioMedicine Online*. 2024;49(2):103855.
82. Laird M, Glister C, Cheewasopit W, Satchell LS, Bicknell AB, Knight PG. 'Free' inhibin α subunit is expressed by bovine ovarian theca cells and its knockdown suppresses androgen production. *Sci Rep*. 2019;9(1):19793.
83. Chapman C, Cree L, Shelling AN. The genetics of premature ovarian failure: current perspectives.

Int J Womens Health. 2015;7:799-810. doi:10.2147/ijwh.S64024

84. Chand AL, Harrison CA, Shelling AN. Inhibin and premature ovarian failure. *Hum Reprod Update*. 2010;16(1):39-50. doi: 10.1093/humupd/dmp031
85. Christofolini DM, Cordts EB, Santos-Pinheiro F, et al. How polymorphic markers contribute to genetic diseases in different populations? The study of inhibin A for premature ovarian insufficiency. *Einstein (Sao Paulo)*. 2017;15(3):269-272. doi: 10.1590/s1679-45082017ao4052
86. qadir. DKJNMAHA. The impact of INH-alpha (rs12720062 G/A) gene polymorphism and serum vitamin D level on risk of premature ovarian failure. *J Popul Ther Clin Pharmacol*. 2023;30(1):336-343. doi: 10.47750/jptcp.2023.1092
87. Sharma S, Meena K, Tabiyad A, Bhardwaj MK, Jaiswal A, Mehta M. Correlation between prolactin, thyroid, LH, FSH, estradiol and progesterone in the infertile women. *Int J Reprod Contracept Obstet Gynecol*. 2023;12(4):1017-1022.
88. Saxena S, Gupta R, Agarwal L, Srivastava PC, Mallick AK. Correlation of serum thyroid stimulating hormone and prolactin in female infertility-a case control study. *IJOGR*. 2016;3(4):388-392.