

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

Authors: Dayo Rotimi Omotoso, Oluwaseun Iwasokun, Oluwasegun Davis Olatomide, Tolulope Timothy Arogundade

Article type: Review

Received: 15 May 2025

Accepted: 29 June 2025

Published online: 2 September 2025

eISSN: 2544-1361

Eur J Clin Exp Med

doi: 10.15584/ejcem.2025.4.18

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our authors we are providing this early version of the manuscript. The manuscript will undergo copyediting and typesetting. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Harnessing therapeutic potential of vitamins and microelements to mitigate testicular damage caused by drugs or chemical toxins – a review

Dayo Rotimi Omotoso, Oluwaseun Iwasokun, Oluwasegun Davis Olatomide, Tolulope Timothy Arogundade

Department of Human Anatomy, Redeemer's University, Ede, Osun State, Nigeria

Correspondence author: Dayo Rotimi Omotoso, e-mail: omotosod@run.edu.ng

ORCID

DRO: <https://orcid.org/0000-0002-4467-2955>

OI: <https://orcid.org/0009-0004-2604-6303>

ODO: <https://orcid.org/0000-0003-4596-3348>

TTA: <https://orcid.org/0000-0002-0179-6181>

ABSTRACT

Introduction and aim. Exposure to drugs and chemical toxins has been a common cause of structural and functional impairment of the male gonad (or testis), often leading to male reproductive disorder and infertility. Health concerns due to drugs or chemical-induced testicular damage have increased the exploration of potential therapeutic agents including vitamins and microelements. This review summarizes therapeutic role of vitamins and microelements against drugs or chemical toxins in preclinical studies.

Material and methods. Relevant articles published on scientific databases like PubMed, Google Scholar, Scopus, and Web of Science were retrieved and critically reviewed for this study.

Analysis of the literature. The mitigating effect of essential vitamins (such as vitamin B₂, vitamin B₉, vitamin B₁₂, vitamin B₁₇, vitamin C, vitamin E) and microelements (such as zinc and selenium) has been demonstrated on testicular damage caused by exposure to drugs and chemical toxins in preclinical studies and associated with their antioxidant, anti-inflammatory and anti-apoptotic properties. This was further characterized with reparation of testicular histopathology, suppression of testicular oxidative damage, improved sperm parameters, elevated testicular antioxidants and testosterone level, upregulation of steroidogenic gene, inhibition of sperm DNA damage.

Conclusion. Vitamins and microelements exert therapeutic effect against drugs and chemical-induced testicular damage.

Keywords. drugs and chemical toxins, testicular damage, vitamins and microelements

Introduction

The testes are primary reproductive organs (or gonads) in males which are responsible for the gametogenic and endocrine functions.¹ Structurally, the testes are composed of seminiferous tubules and interstitial tissue (containing Leydig cells) that perform the spermatogenic and endocrine functions respectively.² The testes are covered by protective fibrous tissue (tunica albuginea) and suspended by the spermatic cords within a pouch (tunica vaginalis of the scrotum) outside the body, thereby enabling the surrounding temperature to be lower than the core body temperature.³

Essentially, the development of secondary sexual characteristics and male reproductive function are achieved under normal morphological, physiological and hormonal profile of the testicular tissue.⁴ Meanwhile, the exposure of testicular tissue to toxic chemical agents, including chemotherapeutic agents such as antibiotics, anti-cancer, non-steroidal anti-inflammatory drugs (NSAIDs) and anabolic steroids, has been reported to cause damage of the testicular tissue morphology and impairment of testicular functions, leading to infertility.⁵⁻⁷

Testicular damage caused by drugs and chemical toxins is further linked with the interruption of testosterone synthesis, disruption of spermatogenesis process, testicular atrophy and when prolonged, reproductive dysfunction and infertility could happen.^{8,9} The mechanisms of drugs and chemical toxins-induced testicular damage essentially involve reproductive hormonal disruption, activation of inflammatory and apoptotic signaling and induction of oxidative stress.¹⁰ The application of potent therapeutic intervention could reverse testicular damage after acute exposure to drugs or chemical toxins while prolonged exposure leads to irreversible testicular damage, reproductive dysfunction and infertility.¹¹

Furthermore, preclinical studies using experimental model of drugs and chemical toxins-induced testicular damage have been conducted to harness the efficacy of potential therapeutic agents such as vitamins and microelements. The therapeutic effects have been characterized by the modulation of associated mechanisms including the induction of oxidative stress, activation of inflammatory and apoptotic signaling, and disruption of genetic regulation of gametogenic and endocrine functions of the testes.

Aim

This review was aimed at summarizing the current information about the therapeutic potential of selected vitamins (including vitamin B₂, vitamin B₉, vitamin B₁₂, vitamin B₁₇, vitamin C, vitamin E) and microelements (including zinc and selenium) in mitigating drugs and chemical toxins-induced testicular damage in preclinical studies. The review further highlighted the mechanisms of therapeutic activity of selected vitamins and microelements.

Material and methods

The scientific publications that are relevant to the current review study were searched across databases including Google scholar, PubMed, Scopus and Web of Science. The keywords employed in the search included: 'male gonadoprotective effect of vitamins', 'male gonadoprotective effect of microelements',

‘antioxidant effect of vitamins’, ‘anti-inflammatory effect of vitamins’, ‘anti-apoptotic effect of vitamins’, ‘antioxidant effect of microelements’, ‘anti-inflammatory effect of microelements’, and ‘anti-apoptotic effect of microelements’. Further in-depth assessment of the identified articles was conducted to extract relevant information regarding the therapeutic activity of selected vitamins and microelements against testicular damage induced by exposure to drugs or chemical toxins in experimental model. Articles containing findings that are relevant to the review study, published in peer-reviewed journals and in English language were regarded as eligible and included in the review. Conversely, duplicated articles and those containing irrelevant information were excluded from the review.

Analysis of the literature

Selected therapeutic agents possess distinct structural formula (Fig. 1) and exhibit different pharmacodynamics and pharmacokinetics which culminate into deleterious effect on the testicular tissue. Similarly, chemical toxins with diverse structural conformation (Fig. 2) exert toxic effects that lead to the morphological and functional impairment of the testes. However, some essential vitamins and microelements have demonstrated therapeutic activity against drugs and chemical toxins-induced testicular primarily based on their antioxidant, anti-inflammatory and anti-apoptotic effects.




Fig. 1. Structural formula of selected drugs with testicular toxic effect




Fig. 2. Structural formula of selected chemicals with testicular toxic effect

Selected drug-induced testicular damage in preclinical studies

Busulfan-induced testicular damage

Busulfan is a chemotherapeutic agent used to treat cancers with characteristic adverse effects such as testicular damage and possibly leading to reproductive dysfunction. The deleterious effect of busulfan on testicular tissue has been characterized with the induction of oxidative stress and inflammation in the testicular tissue, elevated level nitric oxide (NO) and increased myeloperoxidase activity.¹² Other features of busulfan-induced testicular damage included decreased level of testicular glutathione (GSH), increased level of testicular malondialdehyde (MDA), reduced sperm parameters, destruction of seminiferous tubules and degenerated spermatogenic epithelium, Sertoli cells, and Leydig cells.¹³⁻¹⁵ Busulfan-induced testicular toxicity involved reduced activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), and increased level of tumor necrosis factor alpha (TNF- α).^{16,17}

Cisplatin-induced testicular damage

The application of cisplatin, a common antineoplastic agent used to treat different types of cancer, often causes side effect such as testicular damage. The deleterious effect of cisplatin on the testes has been characterized with oxidative damage and upregulation of inflammatory and apoptotic signaling leading to reproductive dysfunction.^{5,18} Cisplatin-induced testicular damage has been characterized with increased level of oxidative stress marker (MDA) and reduction in sperm parameters (such as count, motility, and morphology), and levels of testicular testosterone and antioxidants (SOD, GSH, CAT) as well as distortion of testicular histoarchitecture.^{19,20} Other associated mechanisms of cisplatin-induced testicular damage included upregulated expressions of TNF- α , NF- κ B and caspase-3 factors.¹⁹⁻²¹

Citalopram-induced testicular damage

Citalopram is a selective serotonin reuptake inhibitor widely prescribed to act as a serotonergic antidepressant, with characteristic toxic effects of different body tissues including the testes via induction of oxidative stress.^{22,23} Citalopram-induced testicular damage is characterized with diminished sperm parameters (count, motility, viability, morphology and chromatin integrity), significantly increased NO and MDA levels and apoptotic markers.²³ Other associated mechanisms associated with citalopram-induced testicular damage included decreased GSH, increased sperm DNA damage disruption of reproductive hormones.^{24,25}

Cyclophosphamide-induced testicular damage

Cyclophosphamide is an alkylating drug that function as anti-cancer agent for treatment of cancer and as immunosuppressive agent for treatment of autoimmune and immune-mediated diseases.²⁶ It has been reported to exhibit testicular damage mainly through the induction of oxidative damage within testicular tissue and interference with the genetic expression associated with testicular functions.²⁷ Cyclophosphamide-induced testicular damage has been characterized in experimental model to involve

shrinkage of seminiferous tubules, decline in cellular components (Leydig and Sertoli cells) of testes, marked reduction of sperm parameters, testosterone level, tissue antioxidants while MDA level was significantly elevated.^{27,28} Moreover, testicular damage due to cyclophosphamide exposure showed deleterious effect on testicular histomorphology and leads to impairment of testicular structural and functional integrity.²⁹

Doxorubicin-induced testicular damage

Doxorubicin is a potent anti-neoplastic agent which constitutes the major component of anti-cancer treatment regimens.³⁰ It has been indicated to cause testicular damage mainly through the induction of oxidative stress, inflammation and apoptosis³¹. Doxorubicin-induced testicular damage has been characterized with prominent distortion of testicular histoarchitecture, testicular atrophy, Leydig cell degeneration, oligospermia, dysregulation of 3 β -hydroxysteroid dehydrogenase (3 β -HSD) and 17 β -hydroxysteroid dehydrogenase (17 β -HSD).³²⁻³⁴ Moreover, other associated mechanisms of doxorubicin-induced testicular damage included reduced testicular antioxidants (SOD, CAT, GPx), elevated level of inflammatory and apoptotic markers (IL-1 β , IL-6, TNF- α , Bax, caspase-3); elevated level of MDA, reactive oxygen species, 8-Hydroxy-deoxyguanosine (8-OHdG), 4-Hydroxynonenal (4-HNE).^{31,35,36}

Methotrexate-induced testicular damage

Methotrexate is a drug commonly used for treating diverse kinds of cancer and as an immunosuppressant to treat autoimmune diseases.³⁷ It has been reported to severe adverse including testicular damage. In preclinical studies, methotrexate-induced testicular damage has been characterized with marked decline of sperm parameters (count and motility), testicular antioxidants (SOD, GSH, CAT, GPx), and testosterone level and prominent testicular histopathological changes.^{38,39} Other mechanisms of methotrexate-induced testicular damage included increased levels of MDA, protein carbonyl, and inflammo-apoptotic factors including NO, TNF- α , NF-kB, IL-6, IL-1 β , caspase-3 and p53, reduced expressions of Bcl-2, IL-10, HO-1, EGF, ERK1/2 factors and steroidogenic genes in the testes.³⁹⁻⁴¹

Valproic acid-induced testicular damage

Valproic acid is an anticonvulsant drug used to treat epilepsy. However, it has been indicated to cause tissue toxicities (including the testes) mainly on the basis of induction of oxidative stress and tissue damage. Valproic acid-induced testicular damage has been characterized with marked decline in testosterone levels, sperm parameters (count and normal morphology) and steroidogenic gene expression, as well as prominent testicular histopathological changes.^{42,43} Moreover, valproic acid exposure to testicular tissue further caused abrogation of autophagy, increased inflammatory (NF-kB) and apoptotic (caspase-3) signaling and downregulation of AMPK factor.^{43,44}

Table 1. Profile of selected drugs and mechanisms of testicular damaging effect

Drug	Molecular formula (molecular weight)	Therapeutic applications	Major mechanisms of testicular damage	References
Busulfan	C ₆ H ₁₄ O ₆ S ₂ (246.30 g/mol)	Anti-cancer drug	- reduced sperm parameters - testicular histopathological changes - reduced SOD, CAT, GSH levels - increased levels of MDA, NO, and myeloperoxidase activity - upregulated TNF-α expression	Abarikwu et al. ¹² Abarikwu et al. ¹³ Ezim and Abarikwu ¹⁴ Abd El-Hay et al. ¹⁵ Moghadam et al. ¹⁶ Abarikwu et al. ¹⁷
Cisplatin	Pt(NH ₃) ₂ Cl ₂ (300.10 g/mol)	Anti-cancer drug	- increased level of MDA - reduced sperm parameters - reduced levels of testosterone, testicular antioxidants (SOD, GSH, CAT) - distortion of testicular histoarchitecture - upregulation of TNF-α, NF-κB, caspase-3	Abdel-Latif et al. ⁵ Soni et al. ¹⁸ Elsayed et al. ¹⁹ Nofal et al. ²⁰ Othman et al. ²¹
Citalopram	C ₂₀ H ₂₁ N ₂ O ₂ (405.35 g/mol)	Anti-depressant drug	- diminished sperm parameters - increased NO, MDA levels - decreased testicular GSH level - increased sperm DNA damage - disruption of reproductive hormones	Kraai and Seifert ²² Moradi et al. ²³ Ilgin et al. ²⁴ Moradi et al. ²⁵
Cyclophosphamide	C ₇ H ₁₅ Cl ₂ N ₂ O ₂ P (261.08 g/mol)	Anti-cancer and treatment of autoimmune diseases	- marked testicular histopathology - reduction of sperm parameters - reduced testosterone level - decreased testicular antioxidants - increased MDA level	Ahlmann and Hempel ²⁶ Ghobadi et al. ²⁷ Shaker Kordedeh et al. ²⁸ Adana et al. ²⁹
Doxorubicin	C ₂₇ H ₂₉ NO ₁₁	Anti-cancer drug	- distortion of testicular histoarchitecture	ohan et al. ³²

	(543.52 g/mol)		- dysregulation of testicular hormones - reduced testicular SOD, CAT, GPx - elevated levels of IL-1β, IL-6, TNF-α - upregulation of Bax, caspase-3 - elevated MDA, 8-OHdG, 4-HNE levels	Alafifi et al. ³³ Sarman and Koca ³⁴ Tektemur et al. ³⁵ Ijaz et al. ³⁶
Methotrexate	C ₂₀ H ₂₂ N ₈ O ₅ (454.40 g/mol)	Anti-cancer or anti-rheumatic drug	- decline of sperm parameters - reduced SOD, GSH, CAT, GPx, - reduced testosterone levels - testicular histopathological changes - increased levels of MDA, PC NO, TNF-α, NF-kB, IL-6, IL-1β, - upregulated caspase-3, p53 expressions - reduced expressions of Bcl-2, IL-10, HO-1, and steroidogenic genes	Akacha et al. ³⁸ Hassanein et al. ³⁹ Rofaeil et al. ⁴⁰ Sarman et al. ⁴¹
Valproic acid	C ₈ H ₁₆ O ₂ (144.21 g/mol)	Anticonvulsant drug	- reduced sperm parameters - decline in testosterone level and steroidogenic gene expression - testicular histopathological changes - increased NF-kB, caspase-3 levels - downregulation of AMPK and autophagy	Conei et al. ⁴² Alsemeh et al. ⁴³ Savran et al. ⁴⁴

Selected chemical toxin-induced testicular damage in preclinical studies

Bisphenol A-induced testicular damage

Bisphenol A is a synthetic compound used to manufacture polycarbonate plastics and epoxy resins. As an endocrine disruptor, it has been implicated in preclinical studies as a potent cause of chemical toxin-induced testicular damage whereby the impairment of reproductive function occurs through the characteristic anti-androgenic effect.⁴⁵ Bisphenol A-induced testicular damage was characterized with marked distortion of testicular morphology, elevated levels of oxidative stress and inflammatory markers and increased TUNEL positive cells indicating apoptosis in testis.⁴⁵⁻⁴⁷ Moreover, bisphenol A-induced testicular damage caused significant decline of sperm parameters (viability, motility and normal morphology) and testicular antioxidants (SOD, CAT, GSH) while MDA, MMP, ROS levels were increased.^{48,49}

Fluoride-induced testicular damage

Fluoride is a common environmental contaminant which causes prominent detrimental effect on body tissues such as the testes leading to male reproductive impairment and thereby constitute public health concern.⁵⁰ Fluoride-induced testicular damage and male reproductive impairment was described to involve molecular pathways including apoptosis, DNA damage, hormonal imbalance, inflammatory response, mitochondrial dysfunction.⁵¹ Fluoride exposure caused prominent histopathological and biochemical changes characterized with poor semen quality, reduced testosterone level due to degeneration of Leydig cells, regression of testicular tissue leading to impaired spermatogenesis and infertility⁵². Other indications of fluoride-induced testicular damage included reduced testicular weight and sperm parameters (count, motility, and viability), increased abnormal sperm morphology, induction of oxidative stress and activation of IL-17A pathway.^{53,54}

Monosodium glutamate-induced testicular damage

Monosodium glutamate, chemical commonly used to enhance food flavor, has been reported to cause damaging effects on body tissue (including testes) especially after prolonged exposure via induction of oxidative stress and cell death. Monosodium glutamate-induced testicular damage has been characterized with significant reduction of testosterone level, elevated MDA level, marked testicular histopathological changes.⁵⁵ Other associated mechanisms of monosodium glutamate-induced testicular damage included decreased testicular antioxidants (SOD) and PCNA, Bcl-2 and Beclin 1 expressions; elevated levels of apoptotic and inflammatory markers (TNF- α , Bax, caspase-3) which in turn inhibited the AMPK/mTOR pathway and autophagy in the testes.^{56,57}

Nicotine-induced testicular damage

Nicotine, a naturally occurring alkaloid commonly used in tobacco production and to stimulate pleasure feeling, exhibits potent adverse effect on body tissues including the testes leading to male reproductive dysfunction and infertility. In preclinical studies, nicotine-induced testicular damage was characterized with

significant reduction in testosterone level, testicular sperm parameters (count, viability, normal morphology, motility), marked testicular histopathological changes.^{58,59} Other mechanisms of nicotine-induced testicular damage included marked reduction in SOD and GSH, significantly increased levels of testicular MDA, NO, TBARS.^{59,60} Furthermore, the exposure to nicotine caused reduction of Nrf-2 expression and destruction of blood-testis barrier (BTB) via induction of ferroptosis in testicular cells.⁶¹

Potassium bromate-induced testicular damage

Potassium bromate, a food additive used in production of dough, fish paste, beer, cheese or fermented beverages, has been reported to cause deleterious effects on body tissues including testes via induction of oxidative damage due to accumulation of free radicals.⁶² Potassium bromate-induced testicular damage has been characterized with increased levels of MDA, ROS and RNS, reduced levels of SOD, CAT and TAC, marked distortion of testicular histomorphology.⁶³⁻⁶⁵ Other mechanisms of potassium bromate-induced testicular damage included significant decline of sperm parameters (count, motility, viability, normal morphology) and reproductive hormones including testosterone, LH, FSH, 3HSD, 17HSD.^{63,65,66}

Phthalate-induced testicular damage

Phthalates such as di-(2-ethylhexyl) phthalate (DEHP), Dibutyl phthalate (DBP), and butyl benzyl phthalate (BBP) which are used in making consumer products and plastics, are known environmental toxicants (endocrine disruptors) that cause testicular toxicity through induction of oxidative stress, endocrine disruption, inflammation, and apoptosis.⁶⁷ Phthalates-induced testicular toxicity has been characterized with prominent testicular histopathological changes, decreased testosterone level, reduced sperm parameters, disrupted expression of steroidogenic genes (StAR, CYP11A1, CYP17A1, 17 β -HSD, CYP19A1).^{68,69}

ONLINE 1

Table 2. Profile of selected chemicals and mechanisms of testicular damaging effect

Chemicals	Molecular/Atomic formula (Molecular/atomic mass)	Common applications	Major mechanisms of testicular damage	References
Bisphenol A	C ₁₅ H ₁₆ O ₂ (228.29 g/mol)	- used to produce polycarbonate plastics and epoxy resins	- impairment of reproductive function - marked distortion of testicular morphology - increased TUNEL positive apoptotic cells - decline of sperm parameters - reduced testicular SOD, CAT, GSH - increased MDA, MMP, ROS levels	Bordbar et al. ⁴⁵ Gules et al. ⁴⁶ Tekin and Celebi ⁴⁷ Rezaee-Tazangi et al. ⁴⁸ Asadi-Fard et al. ⁴⁹
Fluoride	F ⁻ (18.99 g/mol)	- used to build healthy teeth, gums, bones	- prominent testicular histopathology - reduced sperm parameters, testosterone level - regression of testicular tissue leading to impaired spermatogenesis - activation of IL-17A pathway	Chhabra et al. ⁵¹ Patial et al. ⁵² Li et al. ⁵³ Talebi et al. ⁵⁴
Monosodium glutamate	C ₅ H ₈ NNa ₂ O ₄ (169.11 g/mol)	- used as food flavor enhancer	- reduction of testosterone level, - elevated MDA level - testicular histopathological changes - decreased testicular antioxidants (SOD) - downregulation of PCNA, Bcl-2, Beclin-1 - upregulation of TNF- α , Bax, caspase-3 - inhibition of AMPK/mTOR pathway	Luqman et al. ⁵⁵ Gad et al. ⁵⁶ Morsy et al. ⁵⁷
Nicotine	C ₁₀ H ₁₄ N ₂ (162.23 g/mol)	- used to stimulate pleasure feeling	- reduced testosterone level, sperm parameters - testicular histopathological changes - reduced levels of SOD and GSH - increased levels of MDA, NO, TBARS	Ikwuka et al. ⁵⁸ Jalili et al. ⁵⁹ Ashoub et al. ⁶⁰ Zhang et al. ⁶¹

			and ferroptosis	
Potassium bromate	KBrO ₃ (167.00 g/mol)	- used as flour additive, for production of malt barley	- downregulated Nrf-2 expression - increased levels of MDA, ROS, RNS - reduced levels of SOD, CAT, TAC - marked testicular histopathology - decline of sperm parameters - reduced level of testosterone, LH, FSH, 3HSD, 17HSD	El-Deeb et al. ⁶² Mohamed et al. ⁶³ Akinola et al. ⁶⁴ Nwonuma et al. ⁶⁵ Akinola et al. ⁶⁶
Phthalates	C ₈ H ₄ O ₄ ⁻² (164.11 g/mol)	- used to increase the flexibility and softness of plastics	- testicular histopathological changes - decreased testosterone level - reduced sperm parameters - disrupted expression of steroidogenic genes	Mondal et al. ⁶⁷ Tang et al. ⁶⁸ Liu et al. ⁶⁹

ONLINE 1

Application of vitamins as mitigants of testicular damage in preclinical studies

Vitamin B₂ (or riboflavin) – therapeutic effect against testicular damage

Vitamin B₂ is a member of the vitamin B family (water-soluble vitamins) which plays essential role in various biological activities and acts as a precursor to flavin mononucleotide and flavin adenine dinucleotide which perform vital roles in enzymatic reactions.⁷⁰⁻⁷² It is an essential nutrient for human health derived from dietary sources (such as egg, fish, nuts, milk, fruits, vegetables and so on) and function as antioxidant that mitigate lipid peroxidation and reperfusion oxidative injury.^{71,73,74} The therapeutic effect of Vitamin B₂ against fluoride-induced testicular damage has been demonstrated and characterized with reparation of testicular histopathological changes, reduction of testicular ROS and SLC3A2 levels, improved sperm parameters, and inhibition of fluoride-induced ferroptosis in testicular Leydig cells which would result into improved testosterone.⁵³

Vitamin B₉ (or folic acid) - therapeutic effect against testicular damage

Vitamin B₉, a water-soluble vitamin which is the synthetic form of folate that are derived from dietary sources. It has been reported to exhibit antioxidant property (comparable to vitamins C and E) and could protect tissues against some diseases caused by harmful oxidation.^{75,76} In a preclinical study, vitamin B₉ had demonstrated therapeutic effect against nicotine-induced testicular damage and characterized its therapeutic activity with reparation of testicular degenerative changes, elevated testosterone level as well as suppression of inflammatory cytokines (TNF- α , IL-6) and oxidative stress markers.⁷⁷ Moreover, the therapeutic effect of vitamin B₉ against bisphenol A-induced testicular damage has been demonstrated in preclinical study and characterized with improved sperm parameters (sperm count, motility, morphology), increased serum testosterone level, amelioration of testicular histopathological changes, as well as decreased TUNEL-positive apoptotic cells.⁴⁶

Vitamin B₁₂ (or cobalamin) – therapeutic effect against testicular damage

Vitamin B₁₂ is a water-soluble vitamin, one of the eight vitamin B family, which participate in metabolism and function as co-factor for two critical biochemical reactions (conversion of l-methylmalonyl coenzyme A to succinyl coenzyme A, and the formation of methionine by methylation of homocysteine).^{78,79} It is totally derived from dietary sources (such as fish, meat, poultry, eggs, milk) and exists in four isoforms cyanocobalamin, hydroxocobalamin, 5'-deoxyadenosylcobalamin, and methylcobalamin.⁸⁰ It has been suggested that Vitamin B₁₂ exhibit antioxidant effect and may protect against induction of oxidative stress and inflammation.⁸¹ In a preclinical study, the therapeutic effect of vitamin B₁₂ against nicotine-induced testicular damage has been assessed and reportedly characterized with amelioration of testicular degenerative changes and increased testosterone level. Other characteristic mechanisms included suppression of inflammatory markers such as TNF- α , IL-6 and markers of oxidative stress.⁷⁷

Vitamin B₁₇ (or amygdalin) – therapeutic effect against testicular damage

Vitamin B₁₇ is a glycoside nutrient derived from dietary sources such as apples and apricots and reported to exhibit certain therapeutic properties especially anti-tumor effect.⁸² Furthermore, the therapeutic effect of vitamin B₁₇ has also been demonstrated in preclinical study against methotrexate-induced testicular damage and characterized with amelioration of testicular histopathology (atrophy and lesions), improved sperm parameters (sperm count, viability, morphology index, total motility, and progressive motility), increased testosterone level, improved level of testicular antioxidants (SOD, CAT, GSH).⁸³

Vitamin C (or ascorbic acid) – therapeutic effect against testicular damage

Vitamin C is a water-soluble which is abundantly derived from natural sources such as fresh fruits like oranges, lemons, grape, leafy green vegetables and exists in the body in ascorbate form.^{84,85} It is a natural antioxidant with ability to inhibit ROS production and scavenge free radicals thereby abrogating oxidative stress that follows exposure to tissue toxins.⁸²⁻⁸⁴ Accordingly, preclinical studies have explored the therapeutic potential of vitamin C to combat testicular pathologies due to exposure to drugs and chemical toxins.

Therapeutic role of vitamin C, based on its potent antioxidant activity, has been demonstrated against cyclophosphamide-induced testicular damage with satisfactory outcomes. This was characterized with restoration of testosterone level, improved sperm parameters and reparation of testicular histopathological changes.⁸⁹ Similarly, preclinical studies involving experimental models of citalopram-induced testicular damage and cisplatin-induced testicular damage have demonstrated therapeutic effect of vitamin C and characterized with improvement of sperm parameters (count, motility, viability, morphology, and chromatin integrity), testosterone level, amelioration of testicular histopathology, reduced levels of MDA, NO, and TUNEL-positive apoptotic cells.^{25,90}

Vitamin E (or alpha-tocopherol) – therapeutic effect against testicular damage

Vitamin E is a fat-soluble vitamin composed of eight essential compounds including four tocopherols and four tocotrienols which are all made up of a chromane ring with a hydroxyl group and side chain.⁹¹ Alpha-tocopherol is a common component which function as an antioxidant that protects the cellular membranes and lipoproteins from peroxidation activity of free radicals.^{66,92,93} As an antioxidant, the therapeutic potential of vitamin E has been explored in preclinical studies using the experimental model of tissue pathologies including the testes. The therapeutic activity of Vitamin E against cisplatin-induced testicular damage has been demonstrated in a preclinical study and characterized with amelioration of testicular histopathology (atrophy and lesions), improvement of sperm parameters (sperm count, motility, viability) testosterone level, testicular weight and lipid profile.⁹⁰

Furthermore, vitamin E mitigated valproic acid-induced testicular damage, characterized with increased testicular weight, improved testicular histology, elevated sperm parameters and testosterone level, upregulation of steroidogenic gene and induction of autophagy through upregulation of LC3, beclin1 and downregulation of p62 factors.⁴³ The study by Akinola et al.⁶⁴ further demonstrated that vitamin E ameliorated potassium bromate-induced testicular damage with the ameliorative activity characterized by

improved sperm parameters (count, motility, viability and normal morphology), diminished levels of ROS, RNS, increased levels of testicular antioxidants (SOD, CAT, TAC) and hormones (testosterone, FSH and LH, 17HSD).

Application of microelements as mitigants of testicular damage in preclinical studies

Zinc – therapeutic effect against testicular damage

Zinc (Zn) is an essential microelement which participates in several body functions and physiological processes including reproduction. Aside its role in enhancing the body antioxidant system, it also acts to promote the activity of vitamin D, suppress prostaglandin synthesis and participate in immune responses and tissue homeostasis.^{94,95} Zn exhibits antioxidant and anti-inflammatory properties which underscore its role in oxidative metabolism of spermatozoa. In addition, it plays vital role in maintenance of epithelial lining of reproductive tracts and exerts regulatory role during the process of capacitation and acrosome reaction.⁹⁶ Hence, Zn deficiency disrupts spermatogenesis and could result into sperm anomalies and reduced testosterone concentration.⁹⁶ The therapeutic role of Zn, based on its potent antioxidant activity, has been demonstrated against cyclophosphamide-induced testicular damage with satisfactory outcomes. This was characterized with restoration of testosterone level, improved sperm parameters, reparation of testicular histopathological changes, reduced oxidative stress markers, and inhibition of sperm DNA damage.^{89,97} The mechanisms of therapeutic role Zn included the modulation of metallothionein (MT), tesmin and Nrf2 pathways.⁹⁷

Selenium – therapeutic effect against testicular damage

Selenium (Se) is a metalloid which is essential to human health in trace amounts but could be harmful in excessive amount.⁹⁸ As an essential microelement involved in improving antioxidant defense, immune functions, and metabolic homeostasis, its deficiency increased susceptibility to various disorders and accelerate aging process leading to reduction of life expectancy.⁹⁵ Se demonstrated protective effect against testicular toxicity primarily based on its antioxidant and anti-inflammatory properties.¹⁰⁰ Accordingly, Se has been explored as potent therapeutic agent in mitigating testicular damage that results from exposure to drugs and chemicals in preclinical studies. The therapeutic activity of Se, through its nanoparticles, has been further demonstrated against monosodium glutamate-induced testicular damage and characterized with reparation of testicular histopathology, elevated testosterone level, reduced MDA level, increased levels of testicular antioxidants (SOD, CAT and GPx).¹⁰¹ Moreover, Se nanoparticles have exhibited ameliorative effect against cisplatin-induced testicular damage. The antioxidant property of Se underscored the therapeutic role of Se which was characterized with increased levels of testicular antioxidants (SOD, GSH, GPx, CAT), decline of MDA level, improvement of sperm parameters (count, motility and abnormal morphology), amelioration of testicular degenerative changes.¹⁰²




Fig. 3. Schematic summary of mechanisms of drugs and chemical toxins-induced testicular damage and therapeutic effect of selected vitamins and microelements

Table 3. Profile and therapeutic role of selected vitamins against drug or chemical-induced testicular damage

Vitamins (other name)	Molecular formula (molecular weight)	Dietary sources	Therapeutic mechanisms against drug or chemical testicular damage	References
Vitamin B ₂ (riboflavin)	C ₁₇ H ₂₀ N ₄ O ₆ (376.37 g/mol)	meat, eggs, almonds, nuts, spinach, apple, kidney bean, spinach, banana, potato, avocado, tomato, oatmeal	- amelioration of testicular histopathology - reduction of testicular ROS - improved sperm parameters - inhibition of ferroptosis in Leydig cells - increased testosterone level	Li et al. ⁵³
Vitamin B ₉ (folic acid)	C ₁₉ H ₁₉ N ₇ O ₆ (176.124 g/mol)	banana, papaya, orange, green peas, kidney bean, crab, tomato juice, avocado, black- eyed pea, peanuts, spinach, beef liver	- reparation of testicular degeneration - elevated testosterone level - suppression of TNF- α , IL-6 - reduced oxidative stress, apoptotic markers - improved sperm parameters - increased testosterone level	Gules et al. ⁴⁶ Ray et al. ⁷⁷
Vitamin B ₁₂ (cobalamin)	C ₁₉ H ₁₉ N ₇ O ₆ (176.124 g/mol)	liver, oyster, mussel, clam, mackerel, crab, sardines, catfish, cod, low-fat beef, skimmed milk, soy milk, salmon, cheese, eggs	- reparation of testicular histopathology - increased testosterone level - suppression of TNF- α , IL-6 - suppression oxidative stress	Ray et al. ⁷⁷
Vitamin B ₁₇ (amygdalin)	C ₁₉ H ₁₉ N ₇ O ₆ (176.124 g/mol)	millet, apple seed oil, strawberry, raspberry, spinach, cassava, barley, almonds, yam, cashew, sweet potatoes	- amelioration of testicular histopathology - improved sperm parameters - increased testosterone level - increased level of SOD, CAT, GSH	Felemban et al. ⁸³
Vitamin C (ascorbic acid)	C ₆ H ₈ O ₆ (441.4 g/mol)	oranges, tomatoes, guava, potato, tomato juice, broccoli,	- increased testosterone level - improved sperm parameters	Moradi et al. ²⁵ Hajjar et al. ⁸⁹

		mango, lemon, pepper, grape fruits, strawberries	- reparation of testicular histopathology - reduced levels of MDA, NO, - reduced TUNEL-positive apoptotic cells	Rauf et al. ⁹⁰
Vitamin E (alpha-Tocopherol)	C ₂₉ H ₅₀ O ₂ (430.71 g/mol)	almonds, nuts, butter, spinach, vegetable oils, olive oil, cereals, meat, egg, avocado, fruits, wheat, cottonseed oil, palm oil, potato, sunflower seeds	- improved sperm parameters - increased testosterone level - improved testicular histology - upregulation of steroidogenic gene - diminished levels of ROS, RNS - increased levels of SOD, CAT, TAC	Alsemeh et al. ⁴³ Akinola et al. ⁶⁴ Rauf et al. ⁹⁰

Table 4. Profile and therapeutic role of selected microelements against drug or chemical-induced testicular damage

Trace element (symbol)	Atomic number (mass)	Sources	Therapeutic mechanisms against drug or chemical testicular damage	References
Zinc (Zn)	30 (65.41 u)	oysters, pork, fishes, liver, meat, wheat, mollusks, dairy products, oats, dried peas, nuts, cheese	- increased testosterone level - improved sperm parameters - reparation of testicular histopathology - reduced oxidative stress markers - inhibition of sperm DNA damage - upregulation of Nrf2 factor	Hajjar et al. ⁸⁹ Maremanda et al. ⁹⁷
Selenium (Se)	34 (76.96 u)	garlic, nuts, cabbage, rice, potatoes, oats, fishes, eggs, meats, lentils	- amelioration of testicular histopathology - elevated testosterone level - improved sperm parameters - reduced MDA level - increased levels of SOD, CAT,	Huang et al. ¹⁰⁰¹⁰¹ Hamza et al. ¹⁰¹ Keshta et al. ¹⁰²

ONLINE FIRST Eur J Clin Exp Med

Conclusion

The damaging effect of drugs and chemical toxins exposure on testicular tissue has been demonstrated to result from the induction of oxidative stress, activation of inflammatory and apoptotic signaling, and characterized with disruption of genetic regulation of gametogenic and endocrine functions of the testes. Hence, potential therapeutic agents with antioxidant, anti-inflammatory and anti-apoptotic properties such as essential vitamins (like Vitamin B₂, Vitamin B₉, Vitamin B₁₂, Vitamin B₁₇, Vitamin C, and Vitamin E) and microelements (like Zinc and Selenium) have demonstrated therapeutic efficacy against testicular damage caused by exposure to drugs and chemical toxins in preclinical studies.

Declaration

Funding

The study has received no funding.

Author contributions

Conceptualization, D.R.O., O.I.; Methodology, D.R.O.; Software, O.I.; Validation, D.R.O., O.I., O.D.O., T.T.A.; Formal Analysis, D.R.O., O.D.O., T.T.A.; Investigation, D.R.O., O.I.; Resources, D.R.O., O.I.; Data Curation, D.R.O., O.I.; Writing – Original Draft Preparation, O.I.; Writing – Review & Editing, D.R.O., O.D.O., T.T.A.; Visualization, D.R.O., O.I.; Supervision, D.R.O.

Conflicts of interest

The authors declare no conflicts of interest.

Data availability

The data that support the relevant findings to the review have been included in the article.

Ethics approval

Not applicable.

References

1. Sembulingam K, Sembulingam P. Male reproductive system. In: *Essential of Medical Physiology*. 6th ed. Jaypee Brothers Medical Publishers ed. New Delhi; 2012:456-458.
2. Das PK, Mukherjee J, Banerjee D. Functional morphology of the male reproductive system. In: *Textbook of Veterinary Physiology*. Berlin, Germany: Springer; 2023:441-476. doi: 10.1007/978-981-19-9410-4_19

3. Kryger JV. Acute and chronic scrotal swelling. In: *Nelson Pediatric Symptom-Based Diagnosis*. Philadelphia, PA: Elsevier; 2018:330-338. doi: 10.1016/B978-0-323-39956-2.00021-2
4. Li L, Lin W, Wang Z. Hormone regulation in testicular development and function. *Int J Mol Sci*. 2024;25(11):5805. doi: 10.3390/ijms25115805
5. Abdel-Latif R, Fathy M, Anwar HA, Naseem M, Dandekar T, Othman EM. Cisplatin-induced reproductive toxicity and oxidative stress: ameliorative effect of kinetin. *Antioxidants*. 2022;11(5):863. doi: 10.3390/antiox11050863
6. Guo R, Lv J, Xu H, Bai Y, Lu B, Han Y. A systems toxicology approach to explore toxicological mechanisms of fluoroquinolones-induced testis injury. *Ecotoxicol Environ Saf*. 2021;228:113002. doi: 10.1016/j.ecoenv.2021.113002
7. Omotoso DR, Akinola AO, Ehiemere WP. Assessment of histo-reparative effect of methanolic extract of *Camellia sinensis* in a rat model of lead-induced testicular damage. *J Drug Deliv Ther*. 2020;10(4-s):46-52. doi: 10.22270/jddt.v10i4-s.4007
8. Lin H. Novel testicular injury biomarkers. In: Dragunow M, ed. *Drug Discovery Toxicology*. Hoboken, NJ: Wiley; 2016:471-474. doi: 10.1002/9781119053248.ch39
9. Taha SHN, Zaghloul HS, Ali AAER, Rashed LA, Sabry RM, Gaballah IF. Molecular and hormonal changes caused by long-term use of high dose pregabalin on testicular tissue: the role of p38 MAPK, oxidative stress and apoptosis. *Mol Biol Rep*. 2020;47(11):8523-8533. doi: 10.1007/s11033-020-05894-6
10. Dutta S, Sengupta, P, Slama P, Roychoudhury S. Oxidative Stress, Testicular Inflammatory Pathways, and Male Reproduction. *Int J Mol Sci*. 2021;22(18):10043. doi: 10.3390/ijms221810043
11. Gabrielsen JS, Tanrikut C. Chronic exposures and male fertility: the impacts of environment, diet, and drug use on spermatogenesis. *Andrology*. 2016;4(4):6480-661. doi: 10.1111/andr.12198
12. Abarikwu SO, Mgbudom-Okah CJ, Onuah CL. The protective effect of rutin against busulfan-induced testicular damage in adult rats. *Drug Chem Toxicol*. 2022;45(3):1035-1043. doi: 10.1080/01480545.2020.1803905
13. Abarikwu SO, Mgbudom-Okah CJ, Njoku R-CC, Okonkwo CJ, Onuoha CC, Wokoma AFS. Gallic acid ameliorates busulfan-induced testicular toxicity and damage in mature rats. *Drug Chem Toxicol*. 2022;45(4):1881-1890. doi: 10.1080/01480545.2021.1892949
14. Ezim OO, Abarikwu SO. Exogenous ascorbate administration elevates testicular oxidative damage and histological injuries in rats after busulfan treatment. *Andrologia*. 2023;1:5209480. doi: 10.1155/2023/5209480
15. Abd El-Hay RI, Hamed WHE, Mostafa Omar N, Refat El-Bassouny D, Gawish SA. The impact of busulfan on the testicular structure in prepubertal rats: A histological, ultrastructural and

- immunohistochemical study. *Ultrastruct Pathol.* 2023;47(5):424-450. doi: 10.1080/01913123.2023.2234470
16. Moghadam MT, Dadfar R, Khorsandi L. The effects of ozone and melatonin on busulfan-induced testicular damage in mice. *JBRA Assist Repr.* 2021;25(2):176-184. doi: 10.5935/1518-0557.20200081
 17. Abarikwu SO, Njoku RCC, John IG, Amadi BA, Mgbudom-Okah CJ, Onuah CL. Antioxidant and anti-inflammatory protective effects of rutin and kolaviron against busulfan-induced testicular injuries in rats. *Syst Biol Reprod Med.* 2022;68(2):151-161. doi: 10.1080/19396368.2021.1989727
 18. Soni KK, Kim HK, Choi BR, et al. Dose-dependent effects of cisplatin on the severity of testicular injury in Sprague Dawley rats: reactive oxygen species and endoplasmic reticulum stress. *Drug Des Dev Therap.* 2016;10:3959-3968. doi: 10.2147/DDDT.S120014
 19. Elsayed A, Elkomy A, Alkafafy M, et al. Testicular toxicity of cisplatin in rats: ameliorative effect of lycopene and N-acetylcysteine. *Environ Sci Poll Res Int.* 2022;29(16):24077–24084. doi: 10.1007/s11356-021-17736-4
 20. Nofal AE, Okdah YA, Rady MI, Hassaan HZ. Gum Acacia attenuates cisplatin toxic effect spermatogenesis dysfunction and infertility in rats. *Int J Biol Macromolec.* 2023;240:124292. doi: 10.1016/j.ijbiomac.2023.124292
 21. Othman EM, Habib HA, Zahran ME, Amin A, Heeba GH. Mechanistic protective effect of cilostazol in cisplatin-induced testicular damage via regulation of oxidative stress and TNF- α /NF- κ B/caspase-3 pathways. *Int J Mol Sci.* 2023;24(16):12651. doi: 10.3390/ijms241612651
 22. Kraai EP, Seifert SA. Citalopram Overdose: A Fatal Case. *J Med Toxicol.* 2015;11(2):232-236. doi: 10.1007/s13181-014-0441-0
 23. Moradi M, Hashemian MA, Douhandeh E, Peysokhan M, Hashemian AH, Faramarzi A. The protective role of melatonin in citalopram-induced reproductive toxicity via modulating nitro-oxidative stress and apoptosis in male mice. *Reprod Toxicol.* 2023;118:108368. doi: 10.1016/j.reprotox.2023.108368
 24. Ilgin, S, Kilic, G, Baysal, M, et al. Citalopram induces reproductive toxicity in male rats. *Birth Defects Res.* 2017;109(7):475-485. doi: 10.1002/bdr2.1010
 25. Moradi M, Hashemian MA, Fathi M, et al. Utility of vitamin C in ameliorating citalopram-induced testicular toxicity via modulating nitro-oxidative stress and apoptosis in mice. *J Biochem Mol Toxicol.* 2024;38(1):e23543. doi: 10.1002/jbt.23543
 26. Ahlmann M, Hempel G. The effect of cyclophosphamide on the immune system: implications for clinical cancer therapy. *Cancer Chemother Pharmacol.* 2016;78(4):661-671. doi: 10.1007/s00280-016-3152-1
 27. Ghobadi E, Moloudizargari M, Asghari MH, Abdollahi M. The mechanisms of cyclophosphamide-induced testicular toxicity and the protective agents. *Exp Opin Drug Metab Toxicol.* 2017;13(5):525-536. doi: 10.1080/17425255.2017.1277205

28. Shaker Kordedeh Z, Ghorbani S, Ahmadi S, Soleimani Mehranjani M. Silymarin mitigates toxic effects of cyclophosphamide on testicular tissue and sperm parameters in mice. *Reprod Biol.* 2024;24(4):100946. doi: 10.1016/j.repbio.2024.100946
29. Adana MY, Imam A, Bello AA, et al. Oral thymoquinone modulates cyclophosphamide-induced testicular toxicity in adolescent Wistar rats. *Andrologia.* 2022;54(4):e14368. doi: 10.1111/and.14368
30. Kciuk M, Gielecińska A, Mujwar S, et al. Doxorubicin—an agent with multiple mechanisms of anticancer activity. *Cells.* 2023;12(4):659. doi: 10.3390/cells12040659
31. Levi M, Tzabari M, Savion N, Stemmer SM, Shalgi R, Ben-Aharon I. Dexrazoxane exacerbates doxorubicin-induced testicular toxicity. *Reprod.* 2015;150(4):357-366. doi: 10.1530/REP-15-0129
32. Mohan UP, Pichiah T, Iqbal STA, Arunachalam S. Mechanisms of doxorubicin-mediated reproductive toxicity – A review. *Reproductive Toxicology.* 2021;102:80-89. doi: 10.1016/j.reprotox.2021.04.003
33. Alafifi S, Wahdan S, Elsherbiny D, Azab S. Doxorubicin-induced testicular toxicity: possible underlying mechanisms and promising pharmacological treatments in experimental models. *Arch Pharmaceut Sci Ain Shams Univ.* 2022;6(2):196-207. doi: 10.21608/aps.2022.155127.1098
34. Sarman E, Koca HB. Effect of grape seed extract on doxorubicin-induced testicular and epididymal damage in rats. *Human Exp Toxicol.* 2025;44:1-9. doi: 10.1177/09603271251319787
35. Tektemur A, Tektemur NK, Güzel EE. The therapeutic effect of hesperetin on doxorubicin-induced testicular toxicity: Potential roles of the mechanistic target of rapamycin kinase (mTOR) and dynamin-related protein 1 (DRP1). *Toxicol Appl Pharmacol.* 2022;435:115833. doi: 10.1016/j.taap.2021.115833
36. Ijaz MU, Yaqoob S, Hamza A. Apigenin ameliorates doxorubicin prompted testicular damage: biochemical, spermatological and histological based study. *Sci Rep.* 2024;14(1):9049. doi: 10.1038/s41598-024-59392-x
37. Koźmiński P, Halik PK, Chesori R, Gniazdowska E. Overview of Dual-Acting Drug Methotrexate in Different Neurological Diseases, Autoimmune Pathologies and Cancers. *Int J Mol Sci.* 2020;21(10):3483. doi: 10.3390/ijms21103483
38. Akacha A, Badraoui R, Rebai T, Zourgui L. Effect of *Opuntia ficus indica* extract on methotrexate-induced testicular injury: a biochemical, docking and histological study. *J Biomolec Struct Dynam.* 2022;40(10):4341-4351. doi: 10.1080/07391102.2020.1856187
39. Hassanein EHM, Mohamed WR, Hussein RM, Arafa, E.-S. A. Edaravone alleviates methotrexate-induced testicular injury in rats: Implications on inflammation, steroidogenesis, and Akt/p53 signaling. *Int Immunopharmacol.* 2023;117:109969. doi: 10.1016/j.intimp.2023.109969
40. Rofaeil RR, Ibrahim MA, Mohyeldin RH, El-Tahawy NF, Abdelzaher WY. Role of EGF/ERK1/2/HO-1 axis in mediating methotrexate induced testicular damage in rats and the ameliorative effect of

- xanthine oxidase inhibitors. *Immunopharmacol Immunotoxicol.* 2023;45(5):511-520. doi: 10.1080/08923973.2023.2181684
41. Sarman E, Gulle K, Ilhan I. Histochemical, Immunohistochemical, and Biochemical Investigation of the Effect of Resveratrol on Testicular Damage Caused by Methotrexate (MTX). *Reprod Sci.* 2023;30(11):3315-3324. doi: 10.1007/s43032-023-01269-x
 42. Conei D, Rojas M, Santamaría L, Risopatrón J. Protective role of vitamin E in testicular development of mice exposed to valproic acid. *Andrologia.* 2021;53(8):e14140. doi: 10.1111/and.14140
 43. Alsemeh AE, Ahmed MM, Fawzy A, Samy W, Tharwat M, Rezq S. Vitamin E rescues valproic acid-induced testicular injury in rats: Role of autophagy. *Life Sci.* 2022;296:120434. doi: 10.1016/j.lfs.2022.120434
 44. Savran M, Asci H, Armagan İ, et al. Thymoquinone could be protective against valproic acid-induced testicular toxicity by antioxidant and anti-inflammatory mechanisms. *Andrologia.* 2020;52(7) doi: 10.1111/and.13623
 45. Bordbar H, Yahyavi S-S, Noorafshan A, Aliabadi E, Naseh M. Resveratrol ameliorates bisphenol A-induced testicular toxicity in adult male rats: a stereological and functional study. *Basic Clin Androl.* 2023;33(1):1. doi: 10.1186/s12610-022-00174-8
 46. Gules O, Yildiz M, Naseer Z, Tatar M. Effects of folic acid on testicular toxicity induced by bisphenol-A in male Wistar rats. *Biotechnol Histochem.* 2019;94(1):26-35. doi: 10.1080/10520295.2018
 47. Tekin S, Çelebi F. Investigation of the effect of hesperidin on some reproductive parameters in testicular toxicity induced by Bisphenol A. *Andrologia.* 2022;54(10): doi: 10.1111/and.14562
 48. Rezaee-Tazangi F, Zeidooni L, Rafiee Z, et al. Taurine effects on Bisphenol A-induced oxidative stress in the mouse testicular mitochondria and sperm motility. *JBRA Assist Repr.* 2020;24(4):428-435. doi: 10.5935/1518-0557.20200017
 49. Asadi-Fard Y, Soleimani MZ, Khodayar MJ, Khorsandi L, Shirani M, Samimi A. Morin improves Bisphenol-A-induced toxicity in the rat testicular mitochondria and sperms. *JBRA Assist Reprod.* 2023;27(2):174-179. doi: 10.5935/1518-0557.20220010
 50. Zhang D, Li M, Shi T, et al. Protective effects of flavonoids on fluoride-induced testicular DNA damage in mice. *Free Rad Biol Med.* 2025;228:79-92. doi: 10.1016/j.freeradbiomed.2024.12.047
 51. Chhabra V, Meenakshi S, Maity S, et al. Impact of fluoride exposure on reproductive health: Insights into molecular mechanisms and health implications. *Reprod Toxicol.* 2025;135:108907. doi: 10.1016/j.reprotox.2025
 52. Patial B, Khan I, Thakur R, Fishta A. Effects of fluoride toxicity on the male reproductive system: A review. *J Trace Elements Med Biol.* 2024;86:127522. doi: 10.1016/j.jtemb.2024.127522

53. Li X, Yang J, Shi E. Riboflavin alleviates fluoride-induced ferroptosis by IL-17A-independent system Xc/GPX4 pathway and iron metabolism in testicular Leydig cells. *Environ Pollut*. 2024b;344:123332. doi: 10.1016/j.envpol.2024
54. Talebi SF, Seify M, Bhandari RK, Shoorei H, Oskuei SD. Fluoride-induced testicular and ovarian toxicity: evidence from animal studies. *Biol Res*. 2025;58(1):6. doi: 10.1186/s40659-025-00586-6
55. Luqman EM, Ananda AT, Widjiati W, Hendrawan VF. Protective Effect of Apis dorsata Honey on Chronic Monosodium Glutamate-Induced Testicular Toxicity in Mus musculus Mice. *Turk J Pharmaceut Sci*. 2022;19(3):246-250. doi: 10.4274/tjps.galenos.2021.30737
56. Gad FA-M, Farouk SM, Emam MA. Antiapoptotic and antioxidant capacity of phytochemicals from Roselle (Hibiscus sabdariffa) and their potential effects on monosodium glutamate-induced testicular damage in rat. *Environ Sci Pollut Res*. 2021;28(2):2379-2390. doi: 10.1007/s11356-020-10674-7
57. Morsy MM, Hassan HA, Morsi RM, Nafea OE, Farag AI, Ramadan RS. Alogliptin attenuates testicular damage induced by monosodium glutamate in both juvenile and adult male rats by activating autophagy: ROS dependent AMPK/mTOR. *Reproductive Toxicology*. 2025;132:108826. doi: 10.1016/j.reprotox.2024.108826
58. Ikwuka DC, Anyaehie BU, Nwobodo E, Umegbolu EI, Nworgu CC. Ameliorative effects of African walnut on nicotine-induced reproductive toxicity in rat model. *Int J Health Sci (Qassim)*. 2021;15(1):3-8.
59. Jalili C, Mastaneh K, Mona P, Ghanbari A, Mohsen Z, Samira D. Protective effect of gallic acid on nicotine-induced testicular toxicity in mice. *Res Pharmaceut Sci*. 2021;16(4):414-424. doi: 10.4103/1735-5362.319579
60. Ashoub AH, Abdel-Naby DH, Safar MM, El-Ghazaly MA, Kenawy SA. Ameliorative effect of fractionated low-dose gamma radiation in combination with ellagic acid on nicotine-induced hormonal changes and testicular toxicity in rats. *Environ Sci Pollut Res*. 2021;28(18):23287-23300. doi: 10.1007/s11356-020-12334-2
61. Zhang Z, Cheng J, Yang L, et al. The role of ferroptosis mediated by Bmal1/Nrf2 in nicotine-induced injury of BTB integrity. *Free Rad Biol Med*. 2023;200:26-35. doi: 10.1016/j.freeradbiomed.2023.02.024
62. EL-Deeb MEE, Abd-EL-Hafez AAA. Can vitamin C affect the KBrO₃ induced oxidative stress on left ventricular myocardium of adult male albino rats? A histological and immunohistochemical study. *J Microscop Ultrastr*. 2015;3:120-136. doi: 10.1016/j.jmau.2015.03.003
63. Mohamed EAK, Saddek EA. The protective effect of taurine and/or vanillin against renal, testicular, and hematological alterations induced by potassium bromate toxicity in rats. *J Basic Appl Zool*. 2019;80:3. doi: 10.1186/s41936-018-0070-2

64. Akinola AO, Omotoso DR, Oyeyemi AW, Daramola OO. Ameliorative activity of α -tocopherol against potassium bromate-induced reproductive functional and morphological impairments in male rats. *As Pac J Sci Technol*. 2024;29(01).
65. Nwonuma CO, Lawal TA, Acho MA, et al. Protective effects of *Allium cepa*-fortified feed on testicular function alterations by potassium bromate-induced oxidative damage: an in vivo and in silico approach. *Comp Clin Pathol*. 2024;33:453-466. Doi: 10.1007/s00580-024-03566-6
66. Akinola AO, Omotoso DR, Oyeyemi AW, Daramola OO, Emojevwe V. α -Tocopherol Mitigates Adverse Effects of Potassium Bromate on Hematological Parameters and Markers of Hepatic Function in Rat Model. *J Morphol Sci*. 2023;40:89-94.
67. Mondal S, Bandyopadhyay A. Antioxidants in mitigating phthalate-induced male reproductive toxicity: A comprehensive review. *Chemosphere*. 2024;364:143297. doi: 10.1016/j.chemosphere.2024.143297
68. Tang X, Wu S, Shen L, et al. Di-(2-ethylhexyl) phthalate (DEHP)-induced testicular toxicity through Nrf2-mediated Notch1 signaling pathway in Sprague-Dawley rats. *Environ Toxicol*. 33(7):720-728. doi: 10.1002/tox.22559
69. Liu L-L, Yue J-Z, Lu Z-Y, et al. Long-term exposure to the mixture of phthalates induced male reproductive toxicity in rats and the alleviative effects of quercetin. *Toxicol Appl Pharmacol*. 2024;483:116816. doi: 10.1016/j.taap.2024.116816
70. Omotoso D, Joseph D. Comparative therapeutic role of ascorbic acid, α -tocopherol and riboflavin in mitigating hepatotoxicity induced by drugs and chemical toxins – a review. *Eur J Clin Exp Med*. 2025. doi: 10.15584/ejcem.2025.3.4
71. Suwannasom N, Kao I, Pruß A, Georgieva R, Bäuml H. Riboflavin: The Health Benefits of a Forgotten Natural Vitamin. *Int J Mol Sci*. 2020;21(3):950. doi: 10.3390/ijms21030950
72. Murgia C, Dehlia A, Guthridge MA. New insights into the nutritional genomics of adult-onset riboflavin-responsive diseases. *Nutr Metab*. 2023;20(1):42. doi: 10.1186/s12986-023-00764-x
73. Omotoso DR, Amakhabi SA. Histo-reparation of hepatic parenchyma of experimental wistar rats exposed to Ccl4-induced hepatopathy and treated with vitamin B2 and green tea extract. *J Appl Biol Sci*. 2021;15(3):247-256.
74. Saedisomeolia A, Ashoori M. Riboflavin in human health: a review of current evidences. In: Watson RR, ed. *Advances in Food and Nutrition Research*. Vol 83. Cambridge, MA: Academic Press; 2018:57-81. doi: 10.1016/bs.afnr.2017.11.002
75. Gliszczynska-Świgło A. Folates as antioxidants. *Food Chem*. 2007;101(4):1480-1483. doi: 10.1016/j.foodchem.2006.04.022

76. Asbaghi O, Ghanavati M, Ashtary-Larky D, et al. Effects of Folic Acid Supplementation on Oxidative Stress Markers: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Antioxidants*. 2021;10:871. doi: 10.3390/antiox1006087
77. Ray D, Bhattacharjee A, Banerjee O, et al. Folic acid and vitamin B12 ameliorate nicotine-induced testicular toxicity in rats. *Biomedicine*. 2019;39(2):353-368. doi: 10.51248/v39i2.22
78. Rizzo G, Laganà AS. A review of vitamin B12. In: Patel VB, Preedy VR, eds. *Molecular Nutrition*. Cambridge, MA: Academic Press; 2020:105-129. doi: 10.1016/B978-0-12-811907-5.00005-1
79. Moravcová M, Siatka T, Krčmová LK, Matoušová K, Mladěnka P. Biological properties of vitamin B12. *Nutr Res Rev*. 2025;38(1):338-370. doi: 10.1017/S0954422424000210
80. Rakusa ZT, Roskar R, Hickey N, Geremia S. Vitamin B₁₂ in Foods, Food Supplements, and Medicines - A Review of Its Role and Properties with a Focus on Its Stability. *Molecules*. 2023;28(1):240. doi: 10.3390/molecules28010240
81. Olorunnisola OS, Ajayi AF, Okeleji LO, Oladipo AA, Emorioloye JT. Vitamins as antioxidants. *J Food Sci Nutr Res*. 2019;2(3):214-235. doi: 10.26502/jfsnr.2642-11000021
82. Amany MB, Moustafa AAA, Manar HA. Antioxidant activity and defensive role of vitamin B17 to fight cancer. *Sci Res Refract Tech Ceram*. 2020;3(2):1-4. doi: 10.31031/COJTS.2020.03.000556
83. Felemban SG, Aldubayan MA, Alhowail A.H., and Almami, I.S., 2020. Vitamin B17 Ameliorates Methotrexate-Induced Reproductive Toxicity, Oxidative Stress, and Testicular Injury in Male Rats. *Oxid Med Cell Longev*. 2020;2020:4372719. doi: 10.1155/2020/4372719
84. Lykkesfeldt J, Tveden-Nyborg P. The Pharmacokinetics of Vitamin C. *Nutrients*. 2019;11(10): 2412. doi: 10.3390/nu11102412
85. Gęgotek A, Skrzydlewska E. Antioxidative and Anti-Inflammatory Activity of Ascorbic Acid. *Antioxidants*. 2022;11(10);1993. doi: 10.3390/antiox11101993
86. Okwuonu UC, Omotoso DR, Bienonwu EO, Adagbonyin O, Dappa J. Histomorphological profile of liver and kidney tissues of albino Wistar rats following exposure to cadmium-induced damage and ascorbic acid supplementation. *Acad Anat Int*. 2020;6(1):15-19. doi: 10.21276/aanat.2020.6.1.7
87. Omotoso DR, Owonikoko WM, Ehiemere WP. Comparative amelioration of renal histomorphology by ascorbic acid and Camellia sinensis extract in Wistar rats exposed to Lead-induced nephropathy. *Ann Med Res*. 2020;27(8):2161-2165. doi: 10.5455/annalsmedres.2020.02.105
88. Conklin PL, Foyer CH, Hancock RD, Ishikawa T, Smirnoff N. Ascorbic acid metabolism and functions. *J Exp Botany*. 2024;75(9):2599-2603. doi: 10.1093/jxb/erae143
89. Hajjar T, Soleymani F, Vatanchian M. Protective Effect of Vitamin C and Zinc as an Antioxidant Against Chemotherapy-Induced Male Reproductive Toxicity. *J Med Life*. 2020;13(2):138-143. doi: 10.25122/jml-2019-0107

90. Rauf N, Nawaz A, Ullah H, et al. Therapeutic effects of chitosan-embedded vitamin C, E nanoparticles against cisplatin-induced gametogenic and androgenic toxicity in adult male rats. *Environ Sci Poll Res*. 2021;28(40):56319-56332. doi: 10.1007/s11356-021-14516-y
91. Jain P, Singh I, Surana SJ, Shirkhedkar AA. Tocopherols and tocotrienols: the essential vitamin E. In: Puri M, ed. *Bioactive Food Components: Activity in Mechanistic Approach*. 1st ed. Elsevier; 2022:139-154. doi: 10.1016/B978-0-12-823569-0.00009-6
92. Torquato P, Marinelli R, Bartolini D, Galli F. Vitamin E: nutritional aspects. In: Stacchiotti A, Corsetti G, eds. *Molecular Nutrition*. 1st ed. Elsevier; 2020:447-485. doi: 10.1016/B978-0-12-811907-5.00019-1
93. Omotoso DR, Ehiemere WP. Comparative histomorphological assessment of Vitamin E and green tea (*Camellia sinensis*) extract-mediated amelioration of Lead-induced hepatopathy in experimental Wistar rats. *Am J Physiol Biochem Pharmacol*. 2020;10:18-24. doi: 10.5455/ajbpb.20191105104249
94. Kloubert V, Rink L. Zinc as a micronutrient and its preventive role of oxidative damage in cells. *Food Funct*. 2015;6(10):3195-3204. doi: 10.1039/c5fo00630a
95. Rice JM, Zweifach A, Lynes MA. Metallothionein regulates intracellular zinc signaling during CD4(+) T cell activation. *BMC Immunol*. 2016;17:13. doi: 10.1186/s12865-016-0151-2
96. Fallah A, Mohammad-Hasani A, Colagar AH. Zinc is an essential element for male fertility: a review of Zn roles in men's health, germination, sperm quality, and fertilization. *J Reprod Infertil*. 2018;19(2):69-81.
97. Maremanda KP, Khan S, Jena G. Zinc protects cyclophosphamide-induced testicular damage in rat: involvement of metallothionein, tesmin and Nrf2. *Biochem Biophys Res Comm*. 2014;445(3):591-596. doi: 10.1016/j.bbrc.2014.02.055
98. Genchi G, Lauria G, Catalano A, Sinicropi MS, Carocci A. Biological activity of selenium and its impact on human health. *Int J Mol Sci*. 2023;24(3):2633. doi: 10.3390/ijms24032633
99. Bjørklund G, Shanaida M, Lysiuk R, et al. Selenium: an antioxidant with a critical role in anti-aging. *Molecules*. 2022;27(19):6613. doi: 10.3390/molecules27196613
100. Huang H, Li X, Wang Z. Anti-inflammatory effect of selenium on lead-induced testicular inflammation by inhibiting NLRP3 inflammasome activation in chickens. *Theriogenol*. 2020;155:139-149. doi: 10.1016/j.theriogenology.2020.06.01
101. Hamza RZ, Diab AEA. Testicular protective and antioxidant effects of selenium nanoparticles on monosodium glutamate-induced testicular structure alterations in male mice. *Toxicol Rep*. 2020;7:254-260. doi: 10.1016/j.toxrep.2020.01.012
102. Keshta AT, Fathallah AM, Attia YA, Salem EA, Watad SH. Ameliorative effect of selenium nanoparticles on testicular toxicity induced by cisplatin in adult male rats. *Food Chem Toxicol*. 2023;179:113979. doi: 10.1016/j.fct.2023.113979