

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.

Authors: Dayo Rotimi Omotoso, Favour Chinaemerem Egwuatu

Article type: Review

Received: 5 June 2025

Accepted: 18 July 2025

Published online: 23 November 2025

eISSN: 2544-1361

Eur J Clin Exp Med

doi:10.15584/ejcem.2026.1.1

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our authors we are providing this early version of the manuscript. The manuscript will undergo copyediting and typesetting. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Drug-induced nephrotoxicity – a review of therapeutic activity of selenium and zinc in preclinical studies

Dayo Rotimi Omotoso, Favour Chinaemerem Egwuatu

Department of Human Anatomy, Redeemer's University, Ede, Osun State, Nigeria

Corresponding author: Dayo Rotimi Omotoso, e-mail: omotosod@run.edu.ng

ORCID

DRO: <https://orcid.org/0000-0002-4467-2955>

FCE: <https://orcid.org/0009-0002-4937-5922>

ABSTRACT

Introduction and aim. Certain drugs cause nephrotoxicity and renal dysfunction through induction of oxidative stress and activation of inflammatory and apoptotic signaling pathways within the renal tissue. To mitigate drug nephrotoxicity, the therapeutic potential of trace elements such as selenium (Se) and zinc (Zn) has been experimentally explored. The current knowledge and mechanisms are hereby summarized in this review.

Material and methods. This narrative review was carried out through a critical assessment of relevant articles published in scientific databases like Google Scholar, PubMed, Scopus, and Web of Science.

Analysis of the literature. The antioxidant, antiapoptotic and anti-inflammatory properties of Se and Zn culminate in their therapeutic activity against drugs nephrotoxicity. The nephroprotective effect of Se and Zn has been characterized with suppression of renal oxidative stress (reduced malondialdehyde, protein carbonyl and elevated levels of superoxide dismutase, glutathione, glutathione peroxidase, catalase, total antioxidant capacity levels); upregulation of anti-apoptotic and anti-inflammatory markers (Bcl-2, heme oxygenase-1, factor related to nuclear factor erythroid 2; downregulation of pro-apoptotic and pro-inflammatory like inducible nitric oxide synthase, nitric oxide, tumor necrosis factor- α , interleukin-6, nuclear factor kappa light chain enhancer of activated B cells NF- κ B, and Bax, leading to reparation of renal histomorphology and improved renal function (indicated by reduced serum creatinine, urea, BUN levels).

Conclusion. The therapeutic activity of Se and Zn against drugs nephrotoxicity underscores their potential role in the management of nephrotoxicity due to pharmacotherapy.

Keywords. drugs nephrotoxicity, selenium nephroprotection, zinc nephroprotection

Introduction

The kidney is a vital organ that performs several essential functions in the body, particularly related to maintaining body homeostasis.^{1,2} In essence, it plays crucial role in maintaining the overall body health by removing toxic wastes and metabolites, regulating body fluid balance and osmolality, maintaining acid-base equilibrium, secreting hormones, and controlling the arterial pressure.^{1,3,4} Due to its essential functions, the kidney is often exposed to and could bioaccumulate potential toxins (regarded as nephrotoxins or nephrotoxins) which would in turn cause nephrotoxicity.^{3,5} Nephrotoxicity thereby involves the degeneration of renal morphological components and loss of their functionality due to the toxic effects of nephrotoxins including chemotherapeutic agents.^{2,6} Hence, the critical role of the kidney tissue in the removal of xenobiotics (including chemotherapeutic agents) from the body makes it prone to their toxic effects.

Drug-induced kidney toxicity has been characterized with renal histopathological changes such as epithelial necrosis, glomerular congestion, interstitial edema and inflammation, tubular dilatation, which often lead to renal functional impairments and kidney failure.⁷⁻⁹ The associated impairment of renal function has been further characterized by significant alteration of markers of renal function such as glomerular filtration rate, urine output, serum levels of electrolytes, and waste products of protein and muscle metabolism.^{9,10} Essentially, nephrotoxicity is characterized with glomerular damage, inflammation, crystal nephropathy, renal tubular cell toxicity, rhabdomyolysis, and thrombotic microangiopathy.^{11,12} As a risk factor of acute kidney injury and chronic kidney disease, nephrotoxicity poses great public health concern that requires effective therapeutic intervention.

Trace elements are essential elements that are required (in very low concentrations) to play an essential role in many physiological and metabolic processes of the body.¹³ They have been shown to exhibit antioxidant, anti-inflammatory and anti-apoptotic effects which thereby underscore their therapeutic potential against various tissue pathologies.¹⁴ Accordingly, the therapeutic potential of trace elements has been explored in preclinical studies to mitigate or ameliorate drug-induced nephrotoxicity.

Aim

In this review, the aim was to elaborate on the therapeutic activity of trace elements, including selenium and zinc, against drug nephrotoxicity in preclinical studies. Furthermore, the associated mechanisms of therapeutic activity of the selected trace elements were highlighted.

Material and methods

Published articles were sought in multiple scientific databases including Google Scholar, PubMed, Scopus, and Web of Science and assessed to select those that are relevant to the objective of the review. The search keywords included: ‘drug nephrotoxicity’, ‘trace element nephroprotection’, ‘trace elements mitigate drug nephrotoxicity’,

‘antioxidants effect of trace elements’. The literature search was conducted between 1 April and 30 April 2025, and articles selected from preliminary search results were further critically evaluated to identify those that contain relevant findings on the therapeutic role of selenium and zinc against drug-induced nephrotoxicity in preclinical studies. The inclusion criteria included only articles that provided relevant findings, articles published in English and in peer-reviewed journals. Other non-compliant articles were excluded from the review.

Analysis of the literature

Induction of oxidative stress and activation of inflammatory and apoptotic signaling pathways within kidney tissue, following exposure to nephrotoxics (including drugs), have been demonstrated as pivotal cellular mechanisms of the resulting nephrotoxicity.¹⁸ Hence, the antioxidant, anti-inflammatory, and anti-apoptotic properties of selected trace elements (selenium and zinc) underscored the rationale for their therapeutic application to mitigate drug-induced nephrotoxicity in preclinical studies as presented in this review.

Drug-induced nephrotoxicity in preclinical studies

Nephrotoxicity due to drug administration occurs as an adverse effect of pharmacotherapy and commonly presents in the form of acute kidney injury, renal tubular disorders, glomerular damage, and nephrolithiasis and could lead to renal failure.¹⁹ Although, the selected drugs under review have different structural conformation (Fig. 1), preclinical studies have demonstrated their relatively similar nephrotoxic mechanisms (Table 1).




Fig. 1. Molecular structures of selected drugs that cause nephrotoxicity

Acetaminophen (or paracetamol) is a common analgesic and antipyretic agent that has been demonstrated to exhibit nephrotoxicity through the induction of endoplasmic reticulum (ER) stress and apoptosis within kidney tissue.²⁰ Acetaminophen-induced nephrotoxicity has been characterized by distortion of renal histomorphology, elevated levels of kidney injury molecule-1 (KIM-1), interleukin-18 (IL-18), serum blood urea nitrogen (BUN) and creatinine.^{20,21} Acetaminophen exposure further caused upregulation of inducible nitric oxide synthase

(iNOS), PERK, activating transcription factor 6 (ATF6), nuclear factor kappa-light chain-enhancer of activated B cells (NF- κ B), p53, caspases 3 and downregulation of Bcl-2 and Bcl-xL expressions.²⁰ Acetaminophen-induced nephrotoxicity also caused decline of tissue antioxidant levels, including superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx), elevated levels of inflammatory markers (tumor necrosis factor- α (TNF- α), IL-1 β , IL-33) and apoptotic marker (caspase-3).^{21,22} Moreover, acetaminophen exposure caused increase in total oxidant status (TOS), and inhibition of nuclear factor erythroid-related factor 2 (Nrf2) signaling pathway.^{22,23}

Furthermore, cisplatin is a common and effective anticancer drug which has been indicated to cause nephrotoxicity by induction of oxidative stress, inflammatory response and apoptosis of renal tubular cells.²⁴ Cellular mechanisms of cisplatin-induced nephrotoxicity included induction of oxidative stress, ER stress, mitochondrial dysfunction, DNA damage and stress responses such as inflammation, autophagy, cell cycle arrest, senescence, and apoptosis.²⁵ Additionally, exposure to cisplatin caused elevated levels of plasma renal markers including creatinine, urea, uric acid and BUN, reduced levels of SOD, CAT, GPx and glutathione (GSH), renal histopathological changes including renal tubular cell death, vacuolization, vascular congestion.²⁶ Cisplatin-induced nephrotoxicity has been further characterized with increased levels of oxidative stress, inflammatory and apoptotic markers (MDA, NO, iNOS, TNF- α , NF- κ B, IL-1 β , caspase-3 and Bax).^{27,28}

Diclofenac is a non-steroidal anti-inflammatory drug (NSAID), commonly applied as analgesic or anti-inflammatory agent, and has been indicated to cause organ toxicity (like kidney) via increased generation of reactive oxygen species (ROS).²⁹ Diclofenac-mediated nephrotoxicity has been characterized with elevated levels of protein carbonyl (PC), urea, creatinine, uric acid, MDA, hydrogen peroxide, decreased antioxidant enzyme activities (SOD, CAT, GPx) and reduced antioxidant (GSH).^{30, 31} Exposure to Diclofenac exposure further resulted in renal histopathological changes, reduced glomerular filtration rate, up-regulation of inflammatory and apoptotic factors including KIM-1, TNF- α , IL-6, IL-18, NF- κ B, STAT3, Bax, p53, HIF-1 α , caspase-3, and cyclooxygenase-2 (COX-2) while heme oxygenase-1 (HO-1), Nrf2, adenosine 5'-monophosphate-activated protein kinase (AMPK), and sirtuin-1 (SIRT-1) expressions were downregulated.³²⁻³⁴

Doxorubicin is an anthracycline anticancer drug which has been reported to cause a toxic effect on body tissue such as the kidney through ROS production, oxidative stress, apoptosis, inflammation and dysregulated autophagic flow, thereby limiting its clinical application.^{35,36} Doxorubicin-induced nephrotoxicity was characterized by a marked decline of activities of antioxidant enzymes (SOD, CAT, GSH), elevated plasma levels of creatinine, urea, uric acid and renal MDA, cholesterol, calcium and sodium concentrations.³⁷ Moreover, doxorubicin exposure caused renal histopathological changes (including glomerular atrophy, tubular congestion and degeneration, inflammatory cell infiltrations), increased levels of BUN, NO, hydrogen peroxide (H₂O₂), upregulation of NF- κ B, IL-1 β , IL-6, caspase-3 and elevated apoptotic index.^{38,39} Other mechanisms of

doxorubicin-induced nephrotoxicity included decline in renal glutathione reductase (GR) activity and elevated levels of TNF- α and plasma neutrophil gelatinase-associated lipocalin (NGAL).⁴⁰

Furthermore, the application of 5-fluorouracil as an anticancer drug has resulted into serious adverse effects, including hepatotoxicity and nephrotoxicity through induction of oxidative stress, inflammation and apoptosis.^{41,42} 5-fluorouracil-induced nephrotoxicity has been characterized by renal histopathological changes (including tubular congestion, degeneration and atrophic glomeruli), marked increase in serum levels of uric acid, creatinine, urea, NO and MDA, decreased activity of antioxidant enzymes (CAT, SOD, GPx) and GSH level.^{42,43} Other mechanisms of 5-fluorouracil-induced nephrotoxicity included up-regulation of lipocalin-2, KIM-1, increased levels of TNF- α , NF- κ B and IL-6 linked with up-regulated expressions of ERK1 / 2 and VCAM-1, down-regulation of IL-10, Nrf2, HO-1, and FXR factors.^{44,45} In addition, 5-fluorouracil administration further resulted in an increased Bax/Bcl-2 ratio, overexpression of iNOS and upregulation of caspase-3 within renal tissue.⁴⁶

Gentamicin, an aminoglycoside antibiotic widely applied as antibiotic to treat Gram-negative bacterial infections, has been reported to exhibit nephrotoxicity due to induction of oxidative stress and activation of apoptotic and inflammatory signaling pathways.⁴⁷ Exposure to gentamicin in preclinical studies resulted in renal histopathology (such as tubular necrosis and tubulointerstitial inflammation), decreased renal antioxidant enzymes (SOD, GSH, CAT), elevated TOS, oxidative stress index (OSI) MDA, iNOS, NO, TNF- α levels.⁴⁷⁻⁴⁹ Other mechanisms of gentamicin-induced nephrotoxicity included up-regulation of NF- κ B p65, IL-1 β , IL-6, IL-18, p38-MAPK, NGAL, KIM-1, caspase-9, caspase-3 and Bax while Bcl-2, HO-1 and Nrf2 were downregulated within kidney tissue.^{47,50,51}

Methotrexate is an effective anticancer and immunosuppressive drug that exhibits characteristic adverse effect including nephrotoxicity due to associated oxidative damage and inflammatory responses in renal tissue.⁵² Methotrexate-induced nephrotoxicity has been characterized by marked distortion of renal histoarchitecture, elevated serum creatinine, BUN, reduced renal CAT, glutathione-S-transferase (GST), GSH, and increased levels of NO, IL-1 β , TNF- α .^{53,54} In preclinical studies, exposure to methotrexate exposure further resulted in impaired mitochondrial biogenesis, reduced SOD and increased MDA levels, upregulation of TLR-4, TNF- α , NF- κ B, IL-6, caspase-3, Bax, beclin-1, LC-3, and down-regulation of Bcl-2, Nrf2, HO-1.^{52,55,56}

Some other chemotherapeutic agents that exhibit nephrotoxicity essentially through the aforementioned mechanisms include cyclophosphamide (an alkylating anticancer drug), vancomycin (a glycopeptide antibiotic drug), tenofovir/lamivudine/efavirenz (a combination antiretroviral drug) and cyclosporine A (an immunosuppressive agent).⁵⁷⁻⁶⁰

Table 1. General profile of selected drugs and mechanisms of nephrotoxicity in preclinical studies

Selected drugs (Applications)	Molecular formula/weight	Experimental model	Dosage/ Route of administration	Mechanisms of nephrotoxicity	References
Acetaminophen (Analgesic, antipyretic agent)	C ₈ H ₉ NO ₂ / 151.16 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 2 g/kg single dose/ oral - 500 mg/kg single dose, intraperitoneal (ip)	renal histopathological changes - elevated levels of markers of oxidative stress, pro-inflammatory and apoptotic factors. - decreased renal function - reduced renal antioxidant levels - downregulation of anti-apoptotic factors	Coban et al. ²⁰ Aktas et al. ²¹ Ozatik et al. ²² Shi et al. ²³
Cisplatin (Anticancer agent)	Pt(NH ₃) ₂ Cl ₂ / 300.10 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 7 mg/kg single dose/ ip - 5 mg/kg single dose / ip	- decline of renal function; - reduced activities of renal antioxidants - distortion of renal histomorphology - increased levels of oxidative stress markers - increased expressions of pro-inflammatory and apoptotic factors	Tang et al. ²⁵ El-Rhman et al. ²⁶ Shinde et al. ²⁷ Qi et al. ²⁸
Diclofenac (Analgesic, antipyretic agent)	C ₁₄ H ₁₁ Cl ₂ NO ₂ / 296.15 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 50 mg/kg daily for 7 days / ip. - 150 mg/kg daily for 6 days / ip	- decreased renal function - increased levels of oxidative stress markers - decreased activities of renal antioxidants - increased levels of pro-inflammatory markers - upregulation of pro-apoptotic factors - downregulation of anti-inflammatory factors	Moradi et al. ³⁰ Karimi-Matloub et al. ³¹ Alorabi et al. ³² Comez et al. ³³ Mansoure et al. ³⁴
Doxorubicin (Anticancer agent)	C ₂₇ H ₂₉ NO ₁₁ / 543.52 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 15 mg/kg single dose/ i.p. - 20 mg/kg single dose/ i.p.	- decrease in levels of renal antioxidants and reduced renal function - increased levels of oxidative stress markers - prominent renal histopathological changes	Ikewuchi et al. ³⁶ Afsar et al. ³⁷ Altinkaynak et al. ³⁸ Hekmat et al. ³⁹

			- 3 mg/kg daily for 6 weeks / ip.	- upregulation of pro-inflammatory and apoptotic signaling	Al Suleimani et al. ⁴⁰
5-Fluorouracil (Anticancer agent)	C ₄ H ₃ FN ₂ O ₂ / 130.08 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 150 mg/kg single dose (day 8) / ip.	- renal histopathological changes - decreased renal function and activity of renal antioxidants - up-regulation of pro-inflammatory and apoptotic factors - downregulation of anti-inflammatory factors	Famurewa et al. ⁴¹ Mansoori et al. ⁴² El-Gendy et al. ⁴³ Althagafy et al. ⁴⁴ Albadrani et al. ⁴⁵ Al-Ghamdi et al. ⁴⁶
Gentamicin (Antibiotic agent)	C ₂₁ H ₄₃ N ₅ O ₇ / 477.60 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 100 mg/kg daily for 8 days / ip - 100 mg/kg daily from day 8 -14 of the 15-day study / ip	- renal histopathological changes - decreased activities of renal antioxidants - elevated levels of oxidative stress markers - up-regulation of inflammoapoptotic factors - down-regulation of anti-apoptotic signaling	Akila et al. ⁴⁷ Abukhalil et al. ⁴⁸ Saeedavi et al. ⁴⁹ Dik et al. ⁵⁰ Nadeem et al. ⁵¹
Methotrexate (Anticancer or antirheumatic agent)	C ₂₀ H ₂₂ N ₈ O ₅ / 454.44 g/mol	Experimental rats (Wistar, Sprague Dawley)	- 20 mg/kg single dose/ i.p.	- distortion of renal histoarchitecture - elevated serum creatinine, BUN levels - reduced levels of renal antioxidants - upregulation of inflammatory and apoptotic factors - down-regulation of anti-apoptotic expressions	Mishriki et al. ⁵³ Morsy et al. ⁵⁴ Wasfey et al. ⁵⁵ Kandemir et al. ⁵⁶

Therapeutic activity of selenium against drug-induced nephrotoxicity

Selenium (Se) is a trace element vital to human health in trace amounts but could exhibit an adverse effect at high concentration.^{61,62} Deficiency of Se has been associated with increased susceptibility of the body to various pathologies.⁶³ Its important role in body antioxidant defense system, metabolic homeostasis, and immune functions underscores its therapeutic potential, which has been widely harnessed to mitigate tissues' toxicity, including the kidney. In a previous study, intraperitoneal (ip) exposure of Se (0.5 or 1 mg/kg) mitigated cyclophosphamide-induced nephrotoxicity and reversed associated mechanisms via increased total antioxidant capacity (TAC), reduced TOS, OSI, serum creatinine and reparation of renal histomorphology.⁵⁷ The oral administration of Se (0.1 mg/kg for 90 days) further caused a protective effect against tenofovir/lamivudine/efavirenz-induced nephrotoxicity, characterized with elevation of renal antioxidant levels (GSH, SOD, GPx, CAT) levels, reduction of the level of MDA, improved renal function (marked by reduced serum creatinine, uric acid, urea) and amelioration of renal histopathological changes.⁵⁸

Furthermore, exposure to Se nanoparticles (0.5, 1 and 2 mg/kg) demonstrated a protective effect against vancomycin nephrotoxicity through a significant decrease in the levels of MDA, iNOS, NO, TNF- α , and KIM-1, increased Bcl-2 and reduced Bax, caspase-3, caspase-9.⁵⁹ Furthermore, daily ip administration of Se (1 mg/kg for 8 days) mitigated gentamicin-induced nephrotoxicity, indicated by reduced serum levels of renal function markers (urea, creatinine), reduced levels of MDA and PC levels, and amelioration of renal histopathological changes.⁶⁴ Furthermore, administration of Se (1.5 mg/kg/day for 5 days, ip) demonstrated protective effect (in combination exposure with vitamin E) against cisplatin nephrotoxicity, characterized with decrease of plasma levels of MDA, urea, creatinine, elevated levels of GSH, GPx, CAT and reparation of renal histopathological changes (Fig. 2).⁶⁵

Therapeutic activity of zinc against drug-induced nephrotoxicity

Zinc (Zn) is an important trace element that participates in several physiological and biochemical processes of the body.⁶⁶ It further participates in the body's antioxidant system, promotes vitamin D, down-regulates prostaglandin synthesis, and enhance immune responses of the body.^{67,68} Essentially, the antioxidant and anti-inflammatory effects of Zn have been explored to mitigate drug-induced nephrotoxicity. In a previous study, the mitigation of cisplatin-induced nephrotoxicity was demonstrated by Zn administration (6 mg / kg, ip) and further characterized by reduced serum urea, creatinine, MDA, TNF- α levels, down-regulation of renal Bax and heat shock proteins.⁶⁹ The protective effect of Zn administration (25 and 50 mg/kg) has been further demonstrated against gentamicin-induced nephrotoxicity and characterized with restoration of renal function indicated by decreased serum urea, creatinine.⁷⁰

Furthermore, the therapeutic activity of Zn supplementation (10 mg/kg/day for 10 days), based on its antioxidant potential, has been demonstrated against cyclosporine A-induced nephrotoxicity in the experimental model and

indicated by reduced serum levels of creatinine, BUN and kidney tissue damage score.⁷¹ Furthermore, Zinc oxide nanoparticles (ZnONPs) have demonstrated a therapeutic effect against doxorubicin-induced nephrotoxicity indicated by elevated levels of renal antioxidants (SOD, GPx, CAT), decline in MDA levels, downregulation of inflammatory markers (IL-6, NF- κ B), improved renal function and improved renal histomorphology.⁷² Additionally, pretreatment with ZnONPs (5 mg/kg ip) exhibited protective effect against cisplatin nephrotoxicity as demonstrated by reduced serum levels of renal function markers (creatinine, BUN), elevated levels of renal antioxidants (SOD, CAT, GR), reduction of renal MDA level, upregulation of anti-inflammatory markers (HO-1 and Nrf2), downregulation of apoptotic factor (Bax) within the kidney tissue.⁷³




Fig. 2. Schematic summary of mechanisms of drug-induced nephrotoxicity and therapeutic activity of selenium and zinc

Table 3. General Profile and therapeutic mechanisms of selected trace elements against drugs nephrotoxicity in preclinical studies

Trace element (symbol/atomic number/mass)	Sources	Biological functions	Experimental model	Treatment regimen	Therapeutic mechanisms against drug-induced nephrotoxicity	References
Selenium (Se/34/76.96 u)	Garlic, nuts, cabbage, rice, potatoes, oats, fishes, eggs, meats, lentils	- Formation of key enzymes, proteins (selenoproteins) - Antioxidant, anti- cancer, anti- inflammatory effect - Immune booster	Experimental rats (Wistar, Sprague Dawley)	- 0.1 mg/kg for 90 days/oral - 1.5 mg/kg/ daily for 5 days/ip	- elevated renal GSH, SOD, GPx, and CAT levels - reduced serum levels of urea, creatinine, and uric acid - decreased levels of TOS, OSI, MDA, PC, - Down-regulation of iNOS, NO, TNF- α and up-regulation of Bcl-2 expressions - amelioration of renal histopathology	Gunes et al. ⁵⁷ Adikwu et al. ⁵⁸ Mehanna et al. ⁵⁹ Bai et al. ⁶¹ Genchi et al. ⁶² Randjelovic et al. ⁶⁴ Aksoy et al. ⁶⁵
Zinc (Zn/30/65.41 u)	Oysters, pork, fish, liver, meat, wheat, mollusks, dairy products, oats, dried peas, nuts, cheese	- Gene regulation - Enzyme cofactor - Wound repair, hair growth - Development of muscle, bone, and cartilage	Experimental rats (Wistar, Sprague Dawley)	- 10 mg/kg/daily for 10 days/oral - 5 mg/kg single dose /ip - 6 mg/kg single dose/ip	- increased renal SOD, GPx, GR, CAT levels - decrease in serum levels of urea, creatinine, BUN - decreased MDA levels - downregulation of IL-6, NF- κ B, TNF- α , TGF- β and upregulation of Bcl-2, HO-1, Nrf2 expressions - amelioration of renal histopathology	Al-Fartusie et al. ⁶⁶ Lahhoba et al. ⁶⁸ Kone et al. ⁷⁰ Choopani et al. ⁷¹ Elgohary et al. ⁷² Barakat et al. ⁷³

Safety concerns and dose-response considerations

Recent studies have reported some concerns regarding selenium exposure at high dosage levels that include adverse metabolic effects of selenoproteins and their potential to stimulate tumor formation.⁷⁴ Similarly, reports have associated excess zinc exposure with clinical symptoms such as anemia, neutropenia, while the deficiency also poses significant health risks.⁷⁵ Therefore, the dosage and duration of exposure to these trace elements require strict regulation both in health and disease.

Conclusion

Exposure to certain drugs mainly results into nephrotoxicity through induction of oxidative stress and activation of inflammatory and apoptotic signaling pathways within the renal tissue. On the other hand, trace elements such as Selenium and zinc have demonstrated therapeutic activity against drug nephrotoxicity in preclinical studies based on their antioxidant, antiapoptotic, and anti-inflammatory properties. The efficacy of Selenium and Zinc as mitigants of drug nephrotoxicity thereby underscores their potential role in the management of nephrotoxicity that occurs during pharmacotherapy.

Declarations

Funding

The research has received no funding.

Author contributions

Conceptualization, D.R.O. and F.C.E.; Methodology, D.R.O.; Software, F.C.E.; Validation, D.R.O. and F.C.E.; Formal Analysis, D.R.O.; Investigation, F.C.E.; Resources, D.R.O.; Data Curation, D.R.O.; Writing – Original Draft Preparation, F.C.E.; Writing – Review & Editing, D.R.O; Supervision, D.R.O.

Conflicts of interest

The authors declare that they have no competing interests.

Data availability

The data supporting the findings of this review study have been included in the article.

Ethics approval

Not applicable.

References

1. Sembulingam K, Sembulingam P. *Essentials of Medical Physiology*. 5th ed. Jaypee Brothers Medical Publishers Ltd; 2010:288.
2. Al-Naimi MS, Rasheed HA, Hussien NR, Al-Kuraishy HM, Al-Gareeb AI. Nephrotoxicity: role and significance of renal biomarkers in the early detection of acute renal injury. *J Adv Pharm Technol Res*. 2019;10(3):95-99. doi:10.4103/japtr.JAPTR_336_18
3. Omotoso DR, Lawal OS, Olatomide OD, Okojie IG. Nephroprotective effect of *Cissampelos owariensis* extract on renal histomorphology of Wistar rats during exposure to carbon tetrachloride-induced nephropathy. *Asia J Biol*. 2019;8(4):1-10. doi:10.9734/AJOB/2019/v8i430071
4. Molaei E, Molaei A, Abedi F, Hayes WA, Karimi G. Nephroprotective activity of natural products against chemical toxicants: the role of Nrf2/ARE signaling pathway. *Food Sci Nutr*. 2021;9(6):3362-3384. doi:10.1002/fsn3.2320
5. Okwuonu UC, Omotoso DR, Bienonwu EO, Adagbonyin O, Dappa J. Histomorphological profile of liver and kidney tissues of albino Wistar rats following exposure to cadmium-induced damage and ascorbic acid supplementation. *Acad Anat Int*. 2020;6(1):15-19. doi:10.21276/aanat.2020.6.1.5
6. Yadav R, Kumar D, Singh J, Jangra A. Environmental toxicants and nephrotoxicity: implications on mechanisms and therapeutic strategies. *Toxicol*. 2024;504:153784. doi:10.1016/j.tox.2024.153784
7. Omotoso DR, Olajumoke JM. Ameliorative effects of ascorbic acid and *Allium sativum* (garlic) ethanol extract on renal parenchyma of gentamicin-induced nephropathic rats. *J Complement Alt Med Res*. 2020;9(4):1-8. doi:10.9734/JOCAMR/2020/V9I430146
8. Omotoso DR, Owonikoko WM, Ehiemere WP. Comparative amelioration of renal histomorphology by ascorbic acid and *Camellia sinensis* extract in Wistar rats exposed to lead-induced nephropathy. *Ann Med Res*. 2020;27(8):2161-2165. doi:10.5455/annalsmedres.2020.02.105
9. Xu M, Xu S, Yi X. A comparative analysis of drug-induced kidney injury adverse reactions between cyclosporine and tacrolimus based on the FAERS database. *BMC Immunol*. 2025;26:35. doi:10.1186/s12865-025-00714-7
10. Pazhayattil GS, Shirali AC. Drug-induced impairment of renal function. *Int J Nephrol Renovasc Dis*. 2014;7:457-468. doi:10.2147/IJNRD.S39747
11. Kim SY, Moon A. Drug-induced nephrotoxicity and its biomarkers. *Biomol Ther (Seoul)*. 2012;20(3):268-272. doi:10.4062/biomolther.2012.20.3.268
12. Al-Naimi MS, Rasheed HA, Hussien NR, Al-Kuraishy HM, Al-Gareeb AI. Nephrotoxicity: Role and significance of renal biomarkers in the early detection of acute renal injury. *J Adv Pharm Technol Res*. 2019;10(3):95-99. doi:10.4103/japtr.JAPTR_336_18

13. Islam MR, Akash S, Jony MH, et al. Exploring the potential function of trace elements in human health: a therapeutic perspective. *Mol Cell Biochem.* 2023;478:2141-2171. doi:10.1007/s11010-022-04638-3
14. Jahankhani K, Taghipour N, Mashhadi Rafiee M, Nikoonezhad M, Mehdizadeh M, Mosaffa N. Therapeutic effect of trace elements on multiple myeloma and mechanisms of cancer process. *Food Chem Toxicol.* 2023;179:113983. doi:10.1016/j.fct.2023.113983
15. Barker T. Vitamins and human health: systematic reviews and original research. *Nutrients.* 2023;15(13):2888. doi:10.3390/nu15132888
16. Zhang FF, Barr SI, McNulty H, Li D, Blumberg JB. Health effects of vitamin and mineral supplements. *BMJ.* 2020;369:m2511. doi:10.1136/bmj.m2511
17. Rai SN, Singh P, Steinbusch HWM, Vamanu E, Ashraf G, Singh MP. The role of vitamins in neurodegenerative disease: an update. *Biomed.* 2021;9(10):1284. doi:10.3390/biomedicines9101284
18. Barnett LMA, Cummings BS. Cellular and molecular mechanisms of kidney toxicity. *Semin Nephrol.* 2019;39(2):141-151. doi:10.1093/toxsci/kfy159
19. Reis AMM. Drug-induced nephrotoxicity. In: Braund R, ed. *Renal Medicine and Clinical Pharmacy.* Vol 1. Cham, Switzerland: Springer; 2020. doi:10.1007/978-3-030-37655-0_6
20. Çoban FK, İnce S, Demirel HH, İslam İ, Aytuğ H. Acetaminophen-induced nephrotoxicity: suppression of apoptosis and endoplasmic reticulum stress using boric acid. *Biol Trace Elem Res.* 2023;201(1):242-249. doi:10.1007/s12011-022-03114-9
21. Aktas Senocak E, Utlu N, Kurt S, Kucukler S, Mehmet F. Sodium pentaborate prevents acetaminophen-induced hepatorenal injury by suppressing oxidative stress, lipid peroxidation, apoptosis, and inflammatory cytokines in rats. *Biol Trace Elem Res.* 2024;202:1164-1173. doi:10.1007/s12011-023-03755-4
22. Ozatik FY, Teksen Y, Kadioglu E, Ozatik O, Bayat Z. Effects of hydrogen sulfide on acetaminophen-induced acute renal toxicity in rats. *Int Urol Nephrol.* 2019;51(4):745-754. doi:10.1007/s11255-018-2053-0
23. Shi J, Peng X, Huang J, Zhang M, Wang Y. Dihydromyricetin alleviated acetaminophen-induced acute kidney injury via nrf2-dependent anti-oxidative and anti-inflammatory effects. *Int J Mol Sci.* 2025;26(5):2365. doi:10.3390/ijms26052365
24. Volarevic V, Djokovic B, Jankovic MG, et al. Molecular mechanisms of cisplatin-induced nephrotoxicity: a balance on the knife edge between renoprotection and tumor toxicity. *J Biomed Sci.* 2019;26:25. doi:10.1186/s12929-019-0518-9
25. Tang C, Livingston MJ, Safirstein R, Dong Z. Cisplatin nephrotoxicity: new insights and therapeutic implications. *Nat Rev Nephrol.* 2023;19:53-72. doi:10.1038/s41581-022-00631-7

26. El-Rhman RH, El-Naga RN, Gad AM, Tadros MG, Hassaneen SK. Dibenzazepine attenuates against cisplatin-induced nephrotoxicity in rats: involvement of NOTCH pathway. *Front Pharmacol.* 2020;11:567852. doi:10.3389/fphar.2020.567852
27. Shinde SD, Jain PG, Cheke RS, Surana SJ, Gunjegaonkar SM. Abrogation of cisplatin-induced nephrotoxicity in rats and HEK-293 cell lines by formononetin: in vivo and in vitro study. *Comp Clin Pathol.* 2021;30:617-625. doi:10.1007/s00580-021-03252-x
28. Qi J, Gao L. Linarin protects against cisplatin-induced nephrotoxicity via subsiding proinflammatory and oxidative stress biomarkers in male Wistar rats. *Pharmacognosy Mag.* 2024;0(0). doi:10.1177/09731296241297432
29. Famurewa AC, Akunna GG, Nwafor J, Chukwu OC, Ekeleme-Egedigwe CA, Oluniran JN. Nephroprotective activity of virgin coconut oil on diclofenac-induced oxidative nephrotoxicity is associated with antioxidant and anti-inflammatory effects in rats. *Avicenna J Phytomed.* 2020;10(3):316-324
30. Moradi A, Abolfathi M, Javadian M, et al. Gallic Acid Exerts Nephroprotective, Anti-Oxidative Stress, and Anti-Inflammatory Effects Against Diclofenac-Induced Renal Injury in Male rats. *Arch Med Res.* 2021;52(4):380-388. doi:10.1016/j.arcmed.2020.12.005
31. Karimi-Matloub S, Namavari R, Hatefi-Hesari F, et al. The nephroprotective effect of ellagic acid against diclofenac-induced renal injury in male rats: role of Nrf2/HO-1 and NF- κ B/TNF- α pathways. *Biologia* 2022;77:3633-3643. doi:10.1007/s11756-022-01217-1
32. Alorabi M, Cavalu S, Al-Kuraishy HM, et al. Pentoxifylline and berberine mitigate diclofenac-induced acute nephrotoxicity in male rats via modulation of inflammation and oxidative stress. *Biomed Pharmacother.* 2022;152:113225. doi:10.1016/j.biopha.2022.113225
33. Comez M, Cellat M, Kuzu M, et al. The effect of tyrosol on diclofenac sodium-induced acute nephrotoxicity in rats. *J Biochem Mol Toxicol.* 2024;38(1):e23582. doi:10.1002/jbt.23582
34. Mansoure AN, Elshal M, Helal MG. Inhibitory effect of diacerein on diclofenac-induced acute nephrotoxicity in rats via modulating SIRT1/HIF-1 α /NF- κ B and SIRT1/p53 regulatory axes. *Int Immunopharmacol.* 2024;131:111776. doi:10.1016/j.intimp.2024.111776
35. Lu C, Wei J, Gao C, et al. Molecular signaling pathways in doxorubicin-induced nephrotoxicity and potential therapeutic agents. *Int Immunopharmacol.* 2025;144:113373. doi:10.1016/j.intimp.2024.113373
36. Ikewuchi CC, Ifeanchi MO, Ikewuchi JC. Moderation of doxorubicin-induced nephrotoxicity in Wistar rats by aqueous leaf-extracts of *Chromolaena odorata* and *Tridax procumbens*. *Porto Biomed J.* 2021;6(1):e129. doi:10.1097/j.pbj.0000000000000129
37. Afsar T, Razak S, Almajwal A, Al-Disi D. Doxorubicin-induced alterations in kidney functioning, oxidative stress, DNA damage, and renal tissue morphology; improvement by *Acacia hydasypica*

- tannin-rich ethyl acetate fraction. *Saudi J Biol Sci.* 2020;27(9):2251-2260. doi:10.1016/j.sjbs.2020.07.011
38. Altinkaynak Y, Kural B, Akcan BA, et al. Protective effects of L-theanine against doxorubicin-induced nephrotoxicity in rats. *Biomed Pharmacother.* 2018;108:1524-1534. doi:10.1016/j.biopha.2018.09.171
39. Hekmat AS, Chenari A, Alipanah H. et al. Protective effect of alamandine on doxorubicin-induced nephrotoxicity in rats. *BMC Pharmacol Toxicol.* 2021;22:31 doi:10.1186/s40360-021-00494-x
40. Al Suleimani Y, Al Maskari R, Ali BH, et al. Nephroprotective effects of diminazene on doxorubicin-induced acute kidney injury in rats. *Toxicol Rep.* 2023;11:460-468. doi:10.1016/j.toxrep.2023.11.005
41. Famurewa AC, Aja PM, Balogun ME, et al. Prophylactic administration of naringin prevents anticancer drug 5-fluorouracil-induced hepatorenal toxicity via suppressing lipid peroxidation and oxidative stress in rats. *Pharmacol Res Nat Prod.* 2025;6:100137. doi:10.1016/j.prenap.2024.100137
42. Mansoori R, Kazemi S, Almasi D, et al. Therapeutic benefit of melatonin in 5-fluorouracil-induced renal and hepatic injury. *Basic Clin Pharmacol Toxicol.* 2023;134(3):397-411. doi:10.1111/bcpt.13976
43. El-Gendy HF, El-Bahrawy MM, Mansour DA, et al. Unraveling the potential of *Saccharum officinarum* and *Chlorella vulgaris* towards 5-fluorouracil-induced nephrotoxicity in rats. *Pharmaceutics.* 2024;17(7):885. doi:10.3390/ph17070885
44. Althagafy HS, Hassanein EHM. Morin Mitigates 5-Fluorouracil-Induced Nephrotoxicity by Activating Nrf2/HO-1 and FXR, and Suppressing ERK/VCAM-1 and NF- κ B Pathways. *Int Immunopharmacol.* 2025;148:114092. doi:10.1016/j.intimp.2025.114092
45. Albadrani GM, Altyar AE, Kensara OA, et al. Lycopene alleviates 5-fluorouracil-induced nephrotoxicity by modulating PPAR- γ , Nrf2/HO-1, and NF- κ B/TNF- α /IL-6 signals. *Renal Failure.* 2024;46(2). doi:10.1080/0886022X.2024.2423843
46. Al-Ghamdi AH, Mohamed MZ, Elbadry RM, Fouad AA. Kidney protective effect of sitagliptin in 5-fluorouracil-challenged rats. *Pharmacia.* 2024;71:1-5. doi:10.3897/pharmacia.71.e114441
47. Akila AA, Gad RA, Ewees MGED, Abdul-Hamid M, Abdel-Reheim ES. Clopidogrel protects against gentamicin-induced nephrotoxicity through targeting oxidative stress, apoptosis, and coagulation pathways. *Naunyn Schmiedebergs Arch Pharmacol.* 2025;398:2609-2625. doi:10.1007/s00210-024-03380-5
48. Abukhalil MH, Al-Alami Z, Altaie HAA, et al. Galangin prevents gentamicin-induced nephrotoxicity by modulating oxidative damage, inflammation, and apoptosis in rats. *Naunyn Schmiedebergs Arch Pharmacol.* 2025;398:3717-3729. doi:10.1007/s00210-024-03449-1
49. Saeedavi M, Goudarzi M, Fatemi I, et al. Gentisic acid mitigates gentamicin-induced nephrotoxicity in rats. *Tissue Cell.* 2023;84:102191. doi:10.1016/j.tice.2023.102191

50. Dik B, Hatipoglu D, Ates MB. Potential effects of Resatorvid and alpha lipoic acid on gentamicin-induced nephrotoxicity in rats. *Pharmacol Res Persp*. 2024;12(4):e1222. doi:10.1002/prp2.1222
51. Nadeem RI, Aboutaleb AS, Younis NS, Ahmed HI. Diosmin mitigates gentamicin-induced nephrotoxicity in rats: insights on miR-21 and -155 expression, Nrf2/HO-1 and p38-MAPK/NF- κ B pathways. *Toxics*. 2023;11(1):48. doi:10.3390/toxics11010048
52. Rafique Z, Aabid M, Nadeem H, et al. Nephroprotective Potential of 1,3,4-Oxadiazole Derivative Against Methotrexate-Induced Nephrotoxicity in Rats by Upregulating Nrf2 and Downregulating NF- κ B and TNF- α Signaling Pathways. *J Biochem Mol Toxicol*. 2024;38(12):e70084. doi:10.1002/jbt.70084
53. Mishriki AA, Khalifa AK, Ibrahim DA, et al. Empagliflozin mitigates methotrexate-induced nephrotoxicity in male albino rats: insights on the crosstalk of AMPK/Nrf2 signaling pathway. *Futur J Pharm Sci*. 2024;10:95. doi:10.1186/s43094-024-00669-3
54. Morsy MA, El-Sheikh AAK, Abdel-Hafez SMN, Kandeel M, Abdel-Gaber SA. Paeonol Protects Against Methotrexate-Induced Nephrotoxicity via Upregulation of P-gp Expression and Inhibition of TLR4/NF- κ B Pathway. *Front Pharmacol*. 2022;13: 774387. doi:10.3389/fphar.2022.774387
55. Wasfey EF, Shaaban M, Essam M, et al. Infliximab Ameliorates Methotrexate-Induced Nephrotoxicity in Experimental Rat Model: Impact on Oxidative Stress, Mitochondrial Biogenesis, Apoptotic and Autophagic Machineries. *Cell Biochem Biophys*. 2023;81(4):717-726. doi:10.1007/s12013-023-01168-7
56. Kandemir FM, Kucukler S, Caglayan C, Gur C, Batil AA, Gülçin I. Therapeutic effects of silymarin and naringin on methotrexate-induced nephrotoxicity in rats: Biochemical evaluation of anti-inflammatory, antiapoptotic, and antiautophagic properties. *J Food Biochem*. 2017;41(5):e12398. doi:10.1111/jfbc.12398
57. Gunes S, Sahinturk V, Uslu S, Ayhanci A, Kacar S, Uyar R. Protective effects of selenium on cyclophosphamide-induced oxidative stress and kidney injury. *Biol Trace Elem Res*. 2018;185(1):116-123. doi:10.1007/s12011-017-1231-8
58. Adikwu E, Ezerioha CE, Biradei I. Selenium Protects against Tenofovir/Lamivudine/Efavirenz-Induced Nephrotoxicity in Rats. *J Nat Sci Med*. 5(2):157-162. doi:10.4103/jnsm.jnsm_153_20
59. Mehanna ET, Khalaf SS, Mesbah NM, Abo-Elmatty DM, Hafez MM. Anti-oxidant, anti-apoptotic, and mitochondrial regulatory effects of selenium nanoparticles against vancomycin induced nephrotoxicity in experimental rats. *Life Sci*. 2022;288:120098. doi:10.1016/j.lfs.2021.120098
60. Wu Q, Wang X, Nepovimova E, Wang Y, Yang H, Kuca K. Mechanism of cyclosporine A nephrotoxicity: Oxidative stress, autophagy, and signalings. *Food Chem Toxicol*. 2018;118:889-907. doi:10.1016/j.fct.2018.06.054

61. Bai S, Zhang M, Tang S, et al. Effects and impact of selenium on human health, A review. *Molecules*. 2024;30(1):50. doi:10.3390/molecules30010050
62. Genchi G, Lauria G, Catalano A, Sinicropi MS, Carocci A. biological activity of selenium and its impact on human health. *Int J Mol Sci*. 2023;24(3):2633. doi:10.3390/ijms24032633
63. Bjørklund G, Shanaida M, Lysiuk R, et al. Selenium: an antioxidant with a critical role in anti-aging. *Molecules*. 2022;27(19):6613. doi:10.3390/molecules27196613
64. Randjelovic P, Veljkovic S, Stojiljkovic N, et al. Protective effect of selenium on gentamicin-induced oxidative stress and nephrotoxicity in rats. *Drug Chem Toxicol*. 2011;35(2):141-148. doi:10.3109/01480545.2011.589446
65. Aksoy A, Karaoglu A, Akpolat N, Naziroglu M, Ozturk T, Karagoz ZK. Protective role of selenium and high dose vitamin E against cisplatin - induced nephrotoxicity in rats. *Asian Pac J Cancer Prev*. 2015;16(16):6877-6882. doi:10.7314/apjcp.2015
66. Al-Fartusie FS, Mohssan SN. Essential trace elements and their vital roles in human body. *Ind J Adv Chem Sci*. 2017;5(3):127-136. doi:10.22607/IJACS.2017.503003
67. Kloubert V, Rink L. Zinc as a micronutrient and its preventive role of oxidative damage in cells. *Food Func*. 2015;6:3195-3204. doi:10.1039/c5fo00630a
68. Lahhoba QR, Al-sanafd AE, Mohammed NY, et al. Mineral and trace elements, dietary sources, biological effects, deficiency, and toxicity: A review. *Eurasian Chem Commun*. 2023;5:536-555. doi:10.22034/ecc.2023.381964.1594
69. Tuzcu M, Sahin N, Dogukan A, et al. Protective role of zinc picolinate on cisplatin-induced nephrotoxicity in rats. *J Ren Nutr*. 2010;20(6):398-407. doi:10.1053/j.jrn.2010.04.002
70. Kone SD, Gnahoue G, Yapi FH. Evaluation of preventive effects of zinc, vitamin D and their combination against nephrotoxicity induced by gentamicin in rats. *Eur J Biotechnol Biosci*. 2019;7(2):23-29.
71. Choopani S, Kasaei S, Talebi A, et al. Cyclosporine-A induced nephrotoxicity in male and female rats: Is zinc a suitable protective supplement? *Biomed Res Ther*. 5(12): 2888-2897. doi:10.15419/bmrat.v5i12.507
72. Elgohary A, Metwalli F, Mostafa NY, Reffat M, El-Khawaga OY. Zinc oxide nanoparticles regulate NF-kB expression and restrict inflammation response in doxorubicin-induced kidney injury in rats. *Toxicol Environ Health Sci*. 2023;15:437-448. doi:10.1007/s13530-023-00194-5
73. Barakat LAA, Barakat N, Zakaria MM, Khirallah SM. Protective role of zinc oxide nanoparticles in kidney injury induced by cisplatin in rats. *Life Sci*. 2020;262:118503. doi:10.1016/j.lfs.2020.118503
74. Vinceti M, Filippini T, Jablonska E, Saito Y, Wise LA. Safety of selenium exposure and limitations of selenoprotein maximization: Molecular and epidemiologic perspectives. *Environ Res*. 2022;211:113092. doi:10.1016/j.envres.2022.113092

75. Schoofs H, Schmit J, Rink L. Zinc toxicity: understanding the limits. *Molecules*. 2024;29(13):3130. doi:10.3390/molecules29133130.

ONLINE FIRST Eur J Clin Exp Med