



REVIEW PAPER

Natalia Leksa ¹(ABDGF), David Aebisher ²(BDGF), Dorota Bartusik-Aebisher ³(BDGF)

The role of MRI in the central nervous system

¹ Department of Human Anatomy, Medical College of Rzeszów University, Rzeszów, Poland

² Department of Photomedicine and Physical Chemistry, Medical College of Rzeszów University, Rzeszów, Poland

³ Department of Biochemistry and General Chemistry, Medical College of Rzeszów University, Rzeszów, Poland

ABSTRACT

Introduction. Magnetic Resonance Imaging (MRI) has modified the practice of radiology. MRI is base on safe interaction between radiowaves at a particular frequency and hydrogen nuclei in the body. Metabolic encephalopathies are by definition those disorder of the central nervous system that are not due primarily to structural abnormalities.

Aim. Here we present the ¹H MRI and functional MRI (fMRI) method applied to diagnosis of disorders of the central nervous system.

Material and methods. Analysis of literature and self-research.

Results. We have discussed the major MRI applications in the characteristic of the central nervous system. The relationship between the motion of flowing blood and the representation of the blood on images is complex. This work is an introduction to the basic ideas and techniques of fMRI. Therefore, both, ¹H MRI and functional MRI, methods are used in neuroscience.

Conclusion. Noninvasive MRI and functional MRI are daily diagnostics methods in neurology.

Keywords. ¹H MRI, functional MRI, metabolic brain disfunctions

Introduction

Magnetic Resonance Imaging (MRI) has modified the practice of radiology. MRI is base on safe interaction between radiowaves at a particular frequency and hydrogen nuclei in the body. Metabolic encephalopathies are by definition those disorder of the central nervous system that are not due primarily to structural abnormalities.

Fundamental applications of ¹H Magnetic Resonance Imaging in metabolic brain disfunctions are:

- in herpes simplex encephalitis
- in neurosarcoidosis
- in Sjögren's syndrome
- in toxoplasmosis
- in isolated angiitis of the nervous system

Corresponding author: Dorota Bartusik-Aebisher, e-mail: dbartusik-aebisher@ur.edu.pl

Participation of co-authors: A – Author of the concept and objectives of paper; B – collection of data; C – implementation of research; D – elaborate, analysis and interpretation of data; E – statistical analysis; F – preparation of a manuscript; G – working out the literature; H – obtaining funds

Received: 25.08.2019 | Accepted: 19.12.2019

Publication date: March 2020

In herpes simplex encephalitis

Cerebro spinal fluid (CSF) characteristically shows a lymphocytic pleocytosis with mild or moderate elevation of protein and local normal glucose.¹ CSF analysis is a set of laboratory tests. CSF test includes visual observation of color and clarity, tests for glucose, protein, lactate, lactate dehydrogenase, red blood cell count and white blood cell count. Documentation of temporal lobe involvement may be obtained by ECG, radionuclide brain scan, computed tomography (CT) scan, or Magnetic Resonance Imaging (MRI). The role of each of these metabolities in the diagnosis of herpes simplex virus encephalitis is changing.¹ The ECG has been felt by the most consistently useful diagnostic modality early in the course of infection. Radionuclide and CT scans should be used together since either test may be initially normal. The diagnostic accuracy of MRI as compared to radionuclide and CT has much greater clarity of morphologic details has been shown to provide elegant localizing informations in herps simplex encephalitis and in particular with gadolinium enhancement should provide earlier more precise localization of temporal lobe injury than can be achieved using CT.¹

In neurosarcoidosis

A brain CT scan can detect discrete brain mass lesions, multiple nodules hydrocephalus, diffuse inflammation and areas of periventricular white matter hypodensity.² CT scanning is helpful in monitoring disease activity as judged by the size of a mass lesion or the degree of enhancement of diffuse inflammation.² MRI is very sensitive in detecting of sarcoidosis and is good for evaluating the parasellar area, posterior fossa and spinal cord. MRI has demonstrated hypothalamic lesions not seen on CT scan and has revealed clinically unsuspected disease involving white matter in the periventricular white matter in the periventricular regions. Although both CT scan and MRI scan easily demonstrated hydrocephalus, MRI can better define an obstructing lesion at the level of cerebral aqueduct or fourth ventricle. Furthermore, MRI can access CSF flow at sites of possible obstructing lesion at the level of the cerebral aqueduct. MRI with contrast agent is the preferred technique for the evaluating sarcoidosis. Enhanced images frequently reveal diffuse or focal leptomeningeal disease and can demonstrate inflammation. A normal enhanced MRI scan does not exclude the diagnosis of sarcoidosis. MRI angiography has little to contribute in sarcoidosis.²

In Sjögren's syndrome

Sjögren's syndrome is an autoimmune disease affecting lacrimal and salivary glands causing symptoms of xerophthalmia and xerostomia (sicca syndrome). Diagnosis is based on electencephalography and cerebrospinal fluid analysis are each abnormal in about two-thirds of patients with active central nervous system manifesta-

tion. MRI of the head has proved positive in 75 percent of cases with manifestations of Sjögren's syndrome.³

In toxoplasmosis

The lumbar puncture is not helpful in making the diagnosis of toxoplasmosis. The useful finding is a nonspecific lymphocytic pleocytosis. Imaging of the brain with CT is the most useful test for toxoplasmosis.⁴ MRI appears to be more sensitive than for demonstrating the lesions of cerebral toxoplasmosis.

In isolated angiitis of the nervous system

CT and MRI show multiple lesions in both gray and white matter. Normal results of any of the above tests can be suggestive of angiitis and exclude more common explanations for central nervous system. Angiographic findings in cerebral arteritis are classically described as intermittent narrowing and dilatation of blood vessels.⁵ Angiitis confined to small vessels has been found at postmortem in case with normal angiography.

Functional Magnetic Resonance Imaging

The field of functional magnetic resonance (fMRI) has extended enormously since mid-1990 and is dominated by basic neuroscience.⁶⁻¹¹ fMRI is very interdisciplinary and is providing the area to study a detailed map of the brain activation.¹¹⁻¹⁵ The possibility of measuring changes in brain blood flow associated with neural activity in the brain was known from 1998. fMRI results depend from blood volume, a dot of light on the fMRI, the voxel (the pixel of fMRI screens), captures blood flow in the region of approximately 80,000 neurons and more than 4 million synapses.¹⁵⁻²⁴ This dot of light on the screen is the average measurement of activity over one second in this region. However, neuronal signaling occurs a thousand times faster, in the milliseconds.²⁴⁻³⁴

Conclusion

MRI is used in practically the entire body. However, technological advances enable this technique to be applied in neuroscience.

Acknowledgements

Dorota Bartusik-Aebisher acknowledges support from the National Center of Science NCN (New drug delivery systems-MRI study, Grant OPUS-13 number 2017/25/B/ST4/02481).

References

1. Schroth G, Gawehn J, Thron A, Vallbracht A, Voigt K. Early diagnosis of herpes simplex encephalitis by MRI. *Neurology*. 1987;37(2):179-183.
2. Sherman JL, Stern BJ. Sarcoidosis of the CNS: comparison of unenhanced and enhanced MR images. *AJNR Am J Neuroradiol*. 1990;11(5):915-923.

3. Alexander EL. Central nervous system (CNS) manifestations of primary Sjögren's syndrome: an overview. *Scand J Rheumatol Suppl.* 1986;61:161-165.
4. Levy RM, Bredesen DE, Rosenblum ML. Neurological manifestation of the acquired immunodeficiency syndrome. Experience of UCSF and review of the literature. *J Neurosurg.* 1985;62:475-485.
5. Much JR, Herbst KD, Rapaport SI. Thrombosis in patients with the lupus anticoagulant. *Ann Neurol.* 1983;14:155-168.
6. Wang P, Wang Y, Zhang Q. Intracranial hemorrhage after carotid artery stenting in both anterior and posterior circulation: A case report. *Medicine (Baltimore).* 2018;97(51):e13800.
7. Zhang K, Gao BL, Zhao TY, et al. Hybrid operation to revascularize long-segment occluded internal carotid artery prevent further ischemic events. *Neuroradiology.* 2019;61(2):217-224.
8. Chinnadurai P, Bismuth J. Intraoperative Imaging and Image Fusion for Venous Interventions. *Methodist De-bakey Cardiovasc J.* 2018;14(3):200-207.
9. Warner DS, Schwartz BG, Babygirija R, et al. Thrombolysis After Protamine Reversal of Heparin for Acute Ischemic Stroke After Cardiac Catheterization: Case Report and Literature Review. *Neurologist.* 2018;23(6):194-196.
10. Wei M, Ma Y, Yin L. Endovascular treatment for atherosclerotic stenosis within V3 segment of a hypoplastic vertebral artery: Case report. *Medicine (Baltimore).* 2018;97(42):e12048.
11. Larionov AA, Demin DA, Kulikova EA, et al. Limb-shaking transitory ischaemic attack in a patient with subocclusion of the internal carotid artery. *Angiol Sosud Khir.* 2018;24(3):152-156.
12. Hamada S, Kashiwazaki D, Yamamoto S, Akioka N, Kuwayama N, Kuroda S. Impact of Plaque Composition on Risk of Coronary Artery Diseases in Patients with Carotid Artery Stenosis. *J Stroke Cerebrovasc Dis.* 018;27(12):3599-3604.
13. Li K, Ren M, Meng R, Wang F, Ji X. Dural Arteriovenous Fistula Formation Complicated Cerebral Venous Sinus Stenosis After Venous Sinus Stenting. *World Neurosurg.* 2018;120:400-402.
14. Velasco González A, Stracke P, Nordmeyer H, et al. Low rates of recanalization for wide-necked aneurysms treated with stenting after balloon-assisted coiling: combination of techniques delivers stable and improved results during follow-up. *Neuroradiology.* 2018;60(11):1223-1230.
15. Kim KM, Jeong CU, Shim SG, et al. Efficacy and safety of compound tri-metal stent placement for malignant perihilar biliary obstruction. *Niger J Clin Pract.* 2018; 21(9):1121-1126.
16. Salem MM, Maragkos GA, Enriquez-Marulanda A, et al. Statin Therapy and Diabetes Do Not Affect Aneurysm Occlusion or Clinical Outcomes After Pipeline Embolization Device Treatment: A Preliminary Study. *World Neurosurg.* 2018;120:e525-e532.
17. Iwamoto Y, Kitano T, Matsubara S, Uno M, Yagita Y. In-stent thrombosis after carotid artery stenting in a patient with protein C deficiency. *Neurol Sci.* 2018;39(12):2229-2230.
18. Goroshchenko SA, Petrov AE, Rozhchenko LV, Blagorazumova GP, Vyazgina EM, Ivanov AY. Surgical treatment of large and giant vertebrobasilar aneurysms manifested by brainstem compression symptoms. *Zh Vopr Neirokhir Im N N Burdenko.* 2018;82(4):32-37.
19. Huang KL, Chang TY, Ho MY, et al. The correlation of asymmetrical functional connectivity with cognition and reperfusion in carotid stenosis patients. *Neuroimage Clin.* 2018; 9;20:476-484.
20. Zhang Y, Ma C, Liang S, Yan P, Liang F, Guo F, Jiang C. Morphologic Feature Elongation Can Predict Occlusion Status Following Pipeline Embolization of Intracranial Aneurysms. *World Neurosurg.* 2018;119:e934-e940.
21. Cagnazzo F, Lefevre PH, Mantilla D, et al. Patency of the supraclinoid internal carotid artery branches after flow diversion treatment. A meta-analysis. *J Neuroradiol.* 2019;46(1):9-14.
22. Yamaguchi S, Hamabe J, Horie N, et al. Iatrogenic Removal of the Intima in the Middle Cerebral Artery by a Stent Retriever: A Report of Two Cases. *World Neurosurg.* 2018;118:203-208.
23. Okazaki T, Sakamoto S, Shinagawa K, et al. Detection of in-stent protrusion (ISP) by intravascular ultrasound during carotid stenting: Usefulness of stent-in-stent placement for ISP. *Eur Radiol.* 2019;29(1):77-84.
24. Miura M, Yamada K, Shindo S, et al. Optical Frequency Domain Imaging Evaluation of Progressive Carotid Plaque and In-Stent Restenosis Lesion with Multiple Neovascularizations. *World Neurosurg.* 2018;119:54-57.
25. Ito H, Uchida M, Sase T, et al. Risk Factors of Contralateral Microembolic Infarctions Related to Carotid Artery Stenting. *Neurol Med Chir (Tokyo).* 2018;15;58(7):311-319.
26. Tabibian JH, Baron TH. Endoscopic management of primary sclerosing cholangitis. *Expert Rev Gastroenterol Hepatol.* 2018;12(7):693-703.
27. Cai ZQ, Chai SH, Wei XL, You KZ, Li J, Zhang DM. Comparison of postsurgical clinical sequences between completely embolized and incompletely embolized patients with wide necked intracranial aneurysms treated with stent assisted coil embolization technique: A STROBE-compliant study. *Medicine (Baltimore).* 2018;97(23):e10987.
28. Rosen AC, Soman S, Bhat J, et al. Convergence Analysis of Micro-Lesions (CAML): An approach to mapping of diffuse lesions from carotid revascularization. *Neuroimage Clin.* 2018;18:553-559.
29. Kohara K, Ishikawa T, Kobayashi T, Kawamata T. Retinal artery occlusion during carotid artery stenting with distal embolic protection device. *Neuroradiol J.* 2018;31(5):504-508.
30. Murata K, Yamauchi S, Kaneshiro Y, Urano Y, Yamagata K, Takami T. A novel technique to visualize true lumen in

endovascular treatment of the occlusive carotid dissection and the usefulness of external-internal carotid collateral channel. *Interv Neuroradiol.* 2018;24(5):533-539.

31. Kim T, Kim CH, Kang SH, Ban SP, Kwon OK. Relevance of Antiplatelet Therapy Duration After Stent-Assisted Coil Embolization for Unruptured Intracranial Aneurysms. *World Neurosurg.* 2018;116:e699-e708.

32. van Rooij SB, van Rooij WJ, Peluso JP, Sluzewski M. The Woven EndoBridge (WEB) as primary treatment for unruptured intracranial aneurysms. *Interv Neuroradiol.* 2018;24(5):475-481.

33. Trivelato FP, Ulhôa AC, Rezende MT, Castro-Afonso LH, Abud DG. Recurrence of a totally occluded aneurysm after treatment with a pipeline embolization device. *BMJ Case Rep.* 2018;2018. pii: bcr-2018-013842.

34. Kotsugi M, Takayama K, Myouchin K, et al. Carotid Artery Stenting: Investigation of Plaque Protrusion Incidence and Prognosis. *JACC Cardiovasc Interv.* 2017;10(8):824-831.